

Revised Draft Environmental Impact Statement/  
Environmental Impact Report

# Truckee River Operating Agreement

---

## Water Quality Appendix

California and Nevada

August 2004

United States Department of the Interior  
Bureau of Reclamation  
Fish and Wildlife Service  
Bureau of Indian Affairs

State of California  
Department of Water Resources

# Water Quality Appendix

## Contents

	Page
I. Overview .....	I
II. Hydrology for Water Quality Assessment .....	I
III. Hydrology for Biological Assessment .....	3
IV. Water Quality Model Inputs .....	4
V. Water Quality Model Calibration .....	5
VI. Water Quality Model Verification .....	7
VII. Overview of Water Quality Simulations .....	7
VIII. Current Condition Temperature and Water Quality Simulations .....	8
IX. No Action Alternative Temperature and Water Quality Simulations .....	8
X. Local Water Supply Alternative Temperature and Water Quality Simulations .....	21
XI. TROA Alternative Temperature and Water Quality Simulations .....	22
XII. Simulated Versus Preferred Water Temperatures for Fish .....	22
References .....	36
Attachment: Beneficial Uses of Surface Waters .....	follows page 40

## **WATER QUALITY APPENDIX**

### **I. OVERVIEW**

Recent model results reflect more accurate boundary conditions for the Dynamic Stream Simulation and Assessment Model with temperature (DSSAMt) riverine water quality model which were derived from Truckee River Watershed Analysis Risk Management Framework (WARMF) model results. Model results incorporate future land use changes and therefore nonpoint source loadings to the riverine model. Model results also incorporate more accurate point source loadings to the riverine model. Recent DSSAMt model results are more reliable and tend to coincide with environmental observations due to model improvements as well as better inputs.

Water quality was analyzed in detail using historical data and the DSSAMt water quality model. Historical data is summarized by Bender (1995). Documentation for the water quality model is provided by Brock and Caupp (RCR04-1.0, 2004). The water quality model was calibrated to wet, median, and dry hydrologic conditions (Brock and Caupp, RCR04-2.0, 2004) and verified to dry hydrologic conditions (Brock and Caupp, RCR04-2.0, 2004). A sensitivity analysis was done to identify major variables and model coefficients (Brock and Caupp, RCR04-2.0, 2004). Current condition, No Action Alternative, Local Water Supply Alternative (LWSA), and TROA Alternative were simulated. For support of the fishery analysis, a temperature analysis for extremely wet, median, and extremely dry water years was done for various fish species and life stages of each species for the current condition and each alternative (Brock and Caupp, RCR04-3.0, 2004, RCR04-5.0, 2004, RCR04-7.0, 2004, and RCR04-9.0, 2004). A water quality analysis was done for wet, median, and dry calendar years for the current condition and each alternative (Brock and Caupp, RCR04-4.0, 2004, RCR04-6.0, 2004, RCR04-8.0, 2004, and RCR04-10.0, 2004).

Flow is the most important variable affecting Truckee River water quality. Typically, wet hydrologic conditions provide the best water quality and dry hydrologic conditions the worst. Selection of representative hydrologic years was critical to analyzing water quality.

Flow data were statistically analyzed for use as a water quality indicator and for use in selecting representative hydrologic years. Analyses were based on water years where possible. A water year begins October of the previous year and extends through September. However, calendar year hydrology was required for the water quality assessment to accommodate state standards that are based on the calendar year that extends from January through December.

### **II. HYDROLOGY FOR WATER QUALITY ASSESSMENT**

Water quality was analyzed using both a single variable analysis and a multiple variable analysis. Probability of flow exceedence based on 95 water years of monthly flow from 1901 to 1995 was used for single variable analysis. Probability of flow exceedence indicates the percent of time the annual flow was exceeded historically. Single variable analysis provided



information on the frequency of occurrence of flows. However, to adequately address hourly, daily, weekly, and often seasonal variations, representative flow years within this 95 year period were used for multiple variable analysis of water quality. This dual approach takes into account the effects of flow and also other major variables such as initial conditions, meteorology, and nutrient and organic loadings.

Data availability and both flows reflecting natural local runoff and flows reflecting the effects of storage operations were considered in selecting wet, median, and dry years for multiple variable water quality analysis. Supporting documents may use the term average instead of median. Average does not imply the statistical mean, but rather is a general term to denote central tendency of flows or near median. Supporting documents may also use the term violated instead of exceeded. An exceedence does not imply a violation which is an enforcement term, but rather to denote going outside the range of a desired criteria.

Annual averages of natural local runoff just upstream of Farad, California, without the effects of storage operations were ranked for both calendar and water years 1901 through 1992. Table 1 compares the probability of exceedences for annual averages for wet, median, and dry years for calendar and water year periods. For years 1986 (wet), 1989 (median), and 1992 (dry), flow exceedence statistics based on calendar years are nearly identical to those for water years. This indicates these years represent the intended flow condition.

Table 1.—Probability of flow exceedences

Probability of flow exceedences for natural flow above Farad based on calendar years 1901 through 1992		
Condition	Calendar year	Probability of flow exceedence (%)
Wet	1986	11.8
Median	1989	50.5
Dry	1992	93.5
Probability of flow exceedences for natural flow above Farad based on water years 1901 through 1992		
Condition	Water year	Probability of flow exceedence (%)
Wet	1986	12.9
Median	1989	49.5
Dry	1992	94.6

Analysis of flows including the effects of storage operations indicated similar wet, median, and dry patterns. Table 2 shows flow ranking just downstream from Reno, Nevada, at Vista, Nevada, for the 21 water year periods from 1973 through 1993 which includes the effects of storage operations. This 21-year period includes the flow conditions for the modeled wet, median, and dry years (1986, 1989, and 1992). A ranking of water year 7-day low flows at Farad, California; Vista, Nevada; and Nixon, Nevada (table 2) shows that 1992, the baseline

year, was an extremely dry condition that followed fairly dry conditions and represents near worst case low flow conditions in the Truckee River downstream from Derby Diversion Dam.

Table 2.—7-day low flow and rank for three Truckee River stations

Water year ending September 30	Downstream from reservoirs and upstream of CA/NV State line at Farad, CA		Downstream from Reno, NV, at Vista, NV		Downstream from Derby Diversion Dam at Nixon, NV	
	7-day low flow		7-day low flow		7-day low flow	
	(cfs)	(rank)	(cfs)	(rank)	(cfs)	(rank)
1973	352	14	381	17	44	14
1974	385	19	412	18	123	18
1975	439	20	457	20	130	19
1976	383	18	427	19	145	20
1977	71	6	77	7	23	7
1978	47	2	62	5	20	6
1979	174	9	212	10	29	12
1980	300	12	290	14	33	13
1981	230	10	189	9	24	9
1982	284	11	292	15	52	16
1983	460	21	609	21	464	21
1984	363	15	333	16	64	17
1985	366	16	283	13	27	11
1986	327	13	278	12	44	15
1987	380	17	261	11	25	10
1988	81	7	82	8	20	5
1989	67	5	70	6	23	8
1990	101	8	56	4	11	3
1991	51	3	46	2	8.1	2
1992	57	4	47	3	6.2	1
1993	44	1	44	1	16	4

Note: 7-day low flow is the lowest mean discharge for 7 consecutive days for a water year.

### III. HYDROLOGY FOR BIOLOGICAL ASSESSMENT

For biological environmental assessment of current conditions and alternatives, temperature analysis was done for extremely wet, median, and extremely dry 5-year water-year periods to accommodate storage effects on biology.

Meteorology and flow data for temperature modeling was available from October 1961 through December 1992 and this period was used in selecting 5-year periods of water years from the 92-year record. Table 3 lists the ending water years used for fishery temperature analysis.

Table 3.—Probability of flow exceedences for natural flow upstream of Farad based on 5-year periods for water years 1901 through 1992

Condition	Ending year of 5-year period	Probability of flow exceedence for 5-year period (%) <sup>1</sup>
Extremely wet	1986	1
Median	1966	47
Extremely dry	1992	99

<sup>1</sup> Based on monthly medians of the five water years.

#### IV. WATER QUALITY MODEL INPUTS

A complete description of flow and water quality model inputs and outputs has been documented by Brock and Caupp of Rapid Creek Research, Inc. (RCR04-1.0, 2004 through RCR04-10.0, 2004 ). Table 4 provides a list of water quality supporting documents for the revised DEIS/EIR:

Table 4.—Water quality supporting documents for revised DEIS/EIR

RCR Report No. <sup>1</sup>	Document topic	Approximate number of pages
04-1.0	DSSAMt Program Documentation	110
04-2.0	Calibration to 1986, 1989, 1991, 1992, and 1993	200
04-3.0	Temperature—Current Condition	379
04-4.0	Water quality—Current Condition	709
04-5.0	Temperature—No Action	296
04-6.0	Water quality—No Action	800
04-7.0	Temperature—TROA	296
04-8.0	Water quality—TROA	800
04-9.0s	Temperature-LWSA	296
04-10.0	Water Quality-LWSA	800

<sup>1</sup> Report number indicates year draft was assembled (2004).

Data sources; river and irrigation flow summaries; meteorological data; water temperature boundary conditions; water chemistry boundary conditions; and water quality standards, based on beneficial uses as well as desired fishery criteria, are presented in the above documents as statistics, tables, and graphs.

Hydraulic geometry, flows, meteorology, and water quality constituents are the primary inputs to the water quality model. For the calibration years, initial boundary conditions and timestep boundary conditions were derived from measured data. For simulation of current conditions and alternatives, flow boundary conditions were provided from the monthly Truckee River operations model (operations model). It was assumed that relative percentages of the lumped diversions attributable to each of the active diversions would be the same under current conditions and the alternatives. Water quality boundary conditions were derived from measured data, from statistical correlations, or from the Truckee River Watershed Analysis Risk Management Framework (WARMF) model (Chen and Weintraub, 2002). The WARMF model, a daily watershed loading model, accounts for future land use changes and is a better tool to predict water quality boundary conditions for the hourly DSSAMt riverine water quality model than estimates of future conditions. The 1999 historical land use coverage was used for current conditions. Estimated future land use coverage for the year 2020 was used for future alternatives. Both coverages used the same eleven land use categories. Actual meteorology was used for all years simulated.

Water temperature inputs at the upstream model boundary for current conditions and the alternatives were generated from multiple linear regression equations. Historical air temperatures and flows were used to develop the coefficients for the multiple linear regression equations. Bivariate statistical analysis indicated that a majority of the variation in mean daily water temperature is due to variation in mean daily air temperature (Brock and Caupp, RCR04-3.0, Chapter 11). Flow also accounted for a portion of the variability and was accounted for in the estimation procedure for water temperature at the upstream boundary. Actual meteorology and simulated flows from the operations model were used as inputs to predict water temperature inputs for the model

## **V. WATER QUALITY MODEL CALIBRATION**

The following is a summary of the water quality model calibration. A more detailed description is provided by Brock and Caupp (RCR04-2.0, 2004).

Four calendar years were used for the water quality model calibration (1986-wet, 1989-median, 1992-dry, and 1993-median). The two near median years, 1989 and 1993, differ greatly since only 1993 was operated for cui-ui spawning. Large amounts of water for cui-ui spawning were released from Derby Diversion Dam to Pyramid Lake during 1993. Modeled nutrient and total dissolved solids loadings to Pyramid Lake were also larger during 1993 as shown in table 5.

Calendar year 1993 had the most complete measured data sets, has large flow fluctuations, and has extreme 7-day low flows. Therefore, 1993 covers a range of conditions. Modeled data was compared to measured data with closeness-of-fit statistics and data plots.

Residual error and average error were the primary closeness-of-fit statistics used for comparison. Residual error indicates the average difference between simulated and observed values. Average error is calculated as the absolute value of the residual error summed over a specified period and divided by the number of days. Average error is always higher than

Table 5.—Comparison of calibration simulated annual loadings to Pyramid Lake

Condition	Calendar year	Loadings (kg/yr x 1,000)		
		Phosphorus	Nitrogen	TDS
Dry	1992	1	11	13,724
Median	1989	2	24	18,316
Median (cui-ui)	1993	8	94	33,297
Wet	1986	28	236	101,228

residual error since average error uses the absolute value of both positive and negative differences between modeled and observed values. Residual error is a better indication of calibration adequacy or closeness of fit because positive and negative values can cancel each other.

For temperature during the April to September period, residual error of less than 1 °C is excellent and less than 2 °C is good. Calculated temperature residual error was less than or equal to 2.0 °C, and typically less than 1 °C for all calibration years. For dissolved oxygen, less than 1 mg/L is excellent and less than 2 mg/L is good. Calculated minimum dissolved oxygen residual error was as high as 2.9 mg/L. For 1993, the year with the most complete and accurate data, minimum dissolved oxygen residual error was less than or equal to 0.8 mg/L. The statistics indicate a good dissolved oxygen calibration in most years; however, marginally adequate during extremely low flow conditions. However, only limited dissolved oxygen field data were available for comparison for years 1986, 1989, and 1992. Brock and Caupp summarize average error and residual error by river reach for the calibration years (RCR04-2.0, 2004, Chapter 8). Tables with closeness of fit statistics as well as plots of modeled versus observed data are provided.

Statistical error will reflect inadequacies in data measurement accuracy as well as deficiencies in the modeling calibration. Temperature measurement precision is limited to about 0.2 °C and dissolved oxygen measurement precision to about 0.2 mg/L.

The model calibration was adequate for comparing alternatives to No Action, the baseline condition. The temperature calibration is excellent to good in almost all years and locations. The dissolved oxygen calibration was fair with minimum dissolved oxygen concentrations matching observed data more closely than concentrations of maximum dissolved oxygen. Conservative substances such as total dissolved solids adequately matched observed data. Total nitrogen and total phosphorus concentrations followed expected trends.

Generally, the water quality calibration is better at upstream stations than at downstream stations. Uncertainties in input data sets and “round off” error accumulate in a downstream direction.

Low summer dissolved oxygen sags occur downstream from Reno primarily due to low flows and excessive nutrients, which result in an oxygen demand associated with accumulations of organic material.



## **VI. WATER QUALITY MODEL VERIFICATION**

*Please note that due to limited time, the water quality model verification has not been peer reviewed. The contractor worked toward a calibration that matched or was better than the calibration used for the 1998 draft TROA EIS/EIR. Therefore, the following section is not complete and values or statistics reflect the 1998 draft. Prior to the public draft, this section will be updated to reflect the 2004 verification.*

*The following paragraphs of this section are from the 1998 draft and needs to be updated. After calibration, the water quality model was verified with the independent data set for calendar year 1991, a dry year, and 1995, which was on the moderately wet side. The model temperature verification was good to excellent. Calculated temperature residual error was less than or equal to 1.1 °C. The model dissolved oxygen calibration appeared marginal but was difficult to verify due to minimal measured dissolved oxygen data. Calculated minimum dissolved oxygen residual error was as high as -2.7 mg/L. Brock and Caupp (RCR04-3.0, 2004) provide a detailed analysis of model verification inputs and results.*

*Nine years of hydraulic and temperature data have been modeled when including calibration water quality years, verification water quality years, and fish temperature years, thereby greatly improving the reliability, robustness, and credibility of the DSSAMt model over a wide range of hydrologic and dynamic water quality conditions.*

## **VII. OVERVIEW OF WATER QUALITY SIMULATIONS**

Tables and plots summarizing water quality inputs and outputs for river reaches downstream from Reno were prepared for the current condition simulation (Brock and Caupp, RCR04-4.0, 2004) and for the alternatives (Brock and Caupp, RCR04-6.0, 2004, RCR04-8.0, 2004, and RCR04-10.0). This included information on river and irrigation flows, weather, water temperature boundary conditions, water chemistry boundary conditions, annual plots of daily water temperature and dissolved oxygen, monthly water chemistry versus distance, statistical summaries of simulated water quality, water quality standards, seasonal constituent versus distance, and simulated constituent loads. Simulated constituents such as river temperature, dissolved oxygen, and nutrients provided information on the suitability for aquatic life. In general, temperature, high pH (an indicator of algae), DO, chloride, TDS, and total nitrogen violations occur downstream from Reno in mostly dry hydrologic conditions for all current and future conditions. During dry current and future conditions, dissolved oxygen concentrations are too low for adequate growth and maintenance of aquatic life. However, large non-scouring nutrient-rich flows followed by sudden drops in flow also results in more algal biomass and consequently lower DO. Lower flow results in larger minimum to maximum DO swings.

Truckee River watershed water quality was summarized in the Truckee River Water Quality Settlement Agreement - Federal Water Rights Acquisition Program, Final Environmental Impact Statement (Bureau of Indian Affairs, October 2002.)

Summary tables of the water quality books are shown as Water Quality Appendix DSSAMt tables 1 – 12.

## **VIII. CURRENT CONDITION TEMPERATURE AND WATER QUALITY SIMULATIONS**

The modeled current temperature (Brock and Caupp, RCR04-3.0, 2004) and water quality (Brock and Caupp, RCR04-4.0, 2004) conditions reflect simulated flows from the operations model and should not be compared to historical flows. Current conditions reflect current reservoir operations and current demands on the system

The current conditions simulation indicated that warm temperatures for fish and water quality problems exist minimally in wet, and mostly in median and dry hydrologic conditions. The most severe conditions occur during dry hydrologic conditions. Current system operations do not adequately accommodate water quality.

Temperature and DO were chosen as water quality indicators. Total phosphorus, ortho-phosphorus, total nitrogen, nitrate nitrogen, nitrite nitrogen, ammonia nitrogen, organic nitrogen, pH, total dissolved solids, and chloride were also modeled as supporting information for the water quality analysis.

Modeling and data indicated that water quality tends to be worse during the warm summer and early fall months. However, dissolved oxygen downstream from Derby Diversion Dam is low during the end of April 1989 for current conditions. Low flow and warm air temperatures caused the low DO. This indicates that water quality is highly variable making qualification of water quality conditions by season difficult.

Annual summaries of major model water quality parameters under current conditions are shown in tables DSSAMt 1 through 12 for each modeled reach downstream from Reno for wet (1986), median (1989), and dry (1992) calendar year conditions.

Based on summaries from the draft TROA EIS/EIR (1998), under dry year current conditions, annual summaries indicated that standards for TDS and chloride concentrations to Pyramid Lake, a terminal saline desert lake, are exceeded most of the year. Under dry year current conditions, annual summaries indicated that total nitrogen standards in the reach from Lockwood to Derby Diversion Dam are exceeded about one-third of the year.

## **IX. NO ACTION ALTERNATIVE TEMPERATURE AND WATER QUALITY SIMULATIONS**

No Action represents projected future conditions in the year 2033 without a Truckee River operating agreement or changes in system operation. The major change under No Action is

Table DSSAMt 1. Annual summary of simulated Truckee River quality compared to State of Nevada standards

Alternative: **Current Condition**

Year: **1986 CY**  
Run: **1986 (wet) - RT86C1: Current Cond.**  
Date of Run: **24-Feb-2004**

Constituent	Statistic	Reach					
		Glend-McCar	McCar-Lockw	Lockw-Derby	Derby-Wadsw	Wadsw-Numan	Numan-Pyramid
pH	no. of days > STD	0	0	0	0	29	66
pH	no. of days < STD	0	0	0	0	0	0
Dissolved Oxygen	no. of days < STD (5 or 6 mg/L)	0	0	0	0	0	5
Dissolved Oxygen	no. of days < 5 mg/L*	0	0	0	0	0	5
Chloride	no. of days > STD#	93	0	0	0	0	0
Total Phosphorus	annual average	0.012	0.028	0.039	0.036	0.036	0.036
Ortho Phosphorus	no. of days > STD	0					
Total Nitrogen	annual average	0.177	0.260	0.315	0.287	0.290	0.298
Total Nitrogen	no. of days > STD	1	1	1	1	2	4
Nitrate Nitrogen	no. of days > STD	0	0	0	0	0	0
Nitrite Nitrogen	no. of days > STD	0	1	1	0	0	2
Ammonia Nitrogen (unionized)	no. of days > STD	0	0	0	0	1	2
Total Dissolved Solids	annual average#	70	92	108	100	113	121
Total Dissolved Solids	no. of days > STD#	1	0	0	0	0	0
Temperature	no. of days > STD fish flow not adequate	153	30	32	22	27	37
Temperature	no. of days > STD fish flow adequate ^	na	na	na	36	37	37
Temperature	no. of days > DESIRED fish flow not adequate	164	54	58	65	66	71
Temperature	no. of days > DESIRED fish flow adequate	164	54	58	101	103	104

Notes:

no. of days = number of days of the year when the standard was exceeded within any hour of the day  
^ = determined for period November through June = total of 273 days if flows are adequate for spawning  
na = not applicable  
\* = not a State Standard, included for information purposes only  
# = RMHQ (Recommended to Maintain High Water Quality) used in place of beneficial use standard  
DESIRED = desired condition if flows are adequate to protect juvenile Lahontan cutthroat trout  
STD = Nevada State standard for Truckee River to protect beneficial uses

Table DSSAMt 2. Annual summary of simulated Truckee River quality compared to State of Nevada standards

Alternative: **Current Condition**

Year: **1989 CY**  
 Run: **1989 (median) - RT89C1: Current Cond.**  
 Date of Run: **24-Feb-2004**

Constituent	Statistic	Reach					
		Glend-McCar	McCar-Lockw	Lockw-Derby	Derby-Wadsw	Wadsw-Numan	Numan-Pyramid
pH	no. of days > STD	0	0	34	65	125	127
pH	no. of days < STD	0	0	0	0	0	0
Dissolved Oxygen	no. of days < STD (5 or 6 mg/L)	0	0	0	9	10	25
Dissolved Oxygen	no. of days < 5 mg/L*	0	0	0	6	10	28
Chloride	no. of days > STD#	123	0	0	0	8	11
Total Phosphorus	annual average	0.015	0.036	0.052	0.050	0.047	0.046
Ortho Phosphorus	no. of days > STD	0					
Total Nitrogen	annual average	0.234	0.355	0.425	0.408	0.407	0.409
Total Nitrogen	no. of days > STD	1	1	1	2	3	1
Nitrate Nitrogen	no. of days > STD	0	0	0	0	0	0
Nitrite Nitrogen	no. of days > STD	0	1	1	0	0	0
Ammonia Nitrogen (unionized)	no. of days > STD	0	0	0	0	0	0
Total Dissolved Solids	annual average#	95	121	139	143	204	233
Total Dissolved Solids	no. of days > STD#	60	0	0	0	9	11
Temperature	no. of days > STD fish flow not adequate	155	10	28	25	53	62
Temperature	no. of days > STD fish flow adequate ^	na	na	na	77	80	82
Temperature	no. of days > DESIRED fish flow not adequate	162	48	66	89	99	107
Temperature	no. of days > DESIRED fish flow adequate	162	48	66	149	156	160

Notes:

- no. of days = number of days of the year when the standard was exceeded within any hour of the day
- ^ = determined for period November through June = total of 273 days if flows are adequate for spawning
- na = not applicable
- \* = not a State Standard, included for information purposes only
- # = RMHQ (Recommended to Maintain High Water Quality) used in place of beneficial use standard
- DESIRED = desired condition if flows are adequate to protect juvenile Lahontan cutthroat trout
- STD = Nevada State standard for Truckee River to protect beneficial uses

Table DSSAMt 3. Annual summary of simulated Truckee River quality compared to State of Nevada standards

Alternative: **Current Condition**

Year: **1992 CY**  
Run: **1992 (dry) - RT92C1: Current Cond**  
Date of Run: **24-Feb-2004**

Constituent	Statistic	Reach					
		Glend-McCar	McCar-Lockw	Lockw-Derby	Derby-Wadsw	Wadsw-Numan	Numan-Pyramid
pH	no. of days > STD	0	16	160	94	244	3
pH	no. of days < STD	0	0	0	0	0	0
Dissolved Oxygen	no. of days < STD (5 or 6 mg/L)	0	107	109	149	296	172
Dissolved Oxygen	no. of days < 5 mg/L*	0	107	109	104	253	156
Chloride	no. of days > STD#	366	95	97	363	364	365
Total Phosphorus	annual average	0.023	0.101	0.157	0.147	0.085	0.072
Ortho Phosphorus	no. of days > STD	0					
Total Nitrogen	annual average	0.302	0.640	0.818	0.715	0.569	0.527
Total Nitrogen	no. of days > STD	34	125	122	64	14	17
Nitrate Nitrogen	no. of days > STD	0	0	0	0	0	0
Nitrite Nitrogen	no. of days > STD	0	1	1	0	0	0
Ammonia Nitrogen (unionized)	no. of days > STD	0	0	0	0	0	0
Total Dissolved Solids	annual average#	141	184	211	229	534	667
Total Dissolved Solids	no. of days > STD#	307	2	0	64	366	366
Temperature	no. of days > STD fish flow not adequate	199	96	85	64	88	82
Temperature	no. of days > STD fish flow adequate ^	na	na	na	91	106	101
Temperature	no. of days > DESIRED fish flow not adequate	211	128	123	135	154	147
Temperature	no. of days > DESIRED fish flow adequate	211	128	123	171	189	183

Notes:

- no. of days = number of days of the year when the standard was exceeded within any hour of the day
- ^ = determined for period November through June = total of 273 days if flows are adequate for spawning
- na = not applicable
- \* = not a State Standard, included for information purposes only
- # = RMHQ (Recommended to Maintain High Water Quality) used in place of beneficial use standard
- DESIRED = desired condition if flows are adequate to protect juvenile Lahontan cutthroat trout
- STD = Nevada State standard for Truckee River to protect beneficial uses



Table DSSAMt 4. Annual summary of simulated Truckee River quality compared to State of Nevada standards

Alternative: **No Action**

Year: **1986 CY**  
Run: **1986 (wet) - RT86N1: No Action**  
Date of Run: **24-Feb-2004**

Constituent	Statistic	Reach					
		Glend-McCar	McCar-Lockw	Lockw-Derby	Derby-Wadsw	Wadsw-Numan	Numan-Pyramid
pH	no. of days > STD	0	0	0	15	31	97
pH	no. of days < STD	0	0	0	0	0	0
Dissolved Oxygen	no. of days < STD (5 or 6 mg/L)	0	0	0	0	0	4
Dissolved Oxygen	no. of days < 5 mg/L*	0	0	0	0	0	4
Chloride	no. of days > STD#	63	0	0	0	0	0
Total Phosphorus	annual average	0.013	0.030	0.042	0.039	0.039	0.039
Ortho Phosphorus	no. of days > STD	0					
Total Nitrogen	annual average	0.181	0.275	0.336	0.321	0.325	0.331
Total Nitrogen	no. of days > STD	1	1	1	1	3	4
Nitrate Nitrogen	no. of days > STD	0	0	0	0	0	0
Nitrite Nitrogen	no. of days > STD	0	1	1	0	0	1
Ammonia Nitrogen (unionized)	no. of days > STD	0	0	0	0	1	1
Total Dissolved Solids	annual average#	69	92	108	106	119	126
Total Dissolved Solids	no. of days > STD#	2	0	0	0	0	0
Temperature	no. of days > STD fish flow not adequate	146	25	32	20	27	37
Temperature	no. of days > STD fish flow adequate ^	na	na	na	36	37	39
Temperature	no. of days > DESIRED fish flow not adequate	156	53	58	64	66	72
Temperature	no. of days > DESIRED fish flow adequate	156	53	58	100	103	106

Notes:

- no. of days = number of days of the year when the standard was exceeded within any hour of the day
- ^ = determined for period November through June = total of 273 days if flows are adequate for spawning
- na = not applicable
- \* = not a State Standard, included for information purposes only
- # = RMHQ (Recommended to Maintain High Water Quality) used in place of beneficial use standard
- DESIRED = desired condition if flows are adequate to protect juvenile Lahontan cutthroat trout
- STD = Nevada State standard for Truckee River to protect beneficial uses

Table DSSAMt 5. Annual summary of simulated Truckee River quality compared to State of Nevada standards

Alternative: **No Action**

Year: **1989 CY**  
Run: **1989 (median) - RT89N1: No Action**  
Date of Run: **24-Feb-2004**

Constituent	Statistic	Reach					
		Glend-McCar	McCar-Lockw	Lockw-Derby	Derby-Wadsw	Wadsw-Numan	Numan-Pyramid
pH	no. of days > STD	0	0	29	91	147	138
pH	no. of days < STD	0	0	0	0	0	0
Dissolved Oxygen	no. of days < STD (5 or 6 mg/L)	0	0	0	4	10	26
Dissolved Oxygen	no. of days < 5 mg/L*	0	0	0	0	9	29
Chloride	no. of days > STD#	91	0	0	1	0	10
Total Phosphorus	annual average	0.016	0.040	0.056	0.051	0.048	0.047
Ortho Phosphorus	no. of days > STD	0					
Total Nitrogen	annual average	0.244	0.381	0.463	0.441	0.435	0.435
	no. of days > STD	1	1	1	2	3	1
Nitrate Nitrogen	no. of days > STD	0	0	0	0	0	0
Nitrite Nitrogen	no. of days > STD	0	1	1	0	0	0
Ammonia Nitrogen (unionized)	no. of days > STD	0	0	0	0	0	0
Total Dissolved Solids	annual average#	96	123	143	146	196	227
	no. of days > STD#	60	0	0	0	0	8
Temperature	no. of days > STD fish flow not adequate	160	24	32	22	54	62
	no. of days > STD fish flow adequate ^	na	na	na	77	80	82
	no. of days > DESIRED fish flow not adequate	168	60	70	89	100	107
	no. of days > DESIRED fish flow adequate	168	60	70	149	157	160

Notes:

- no. of days = number of days of the year when the standard was exceeded within any hour of the day
- ^ = determined for period November through June = total of 273 days if flows are adequate for spawning
- na = not applicable
- \* = not a State Standard, included for information purposes only
- # = RMHQ (Recommended to Maintain High Water Quality) used in place of beneficial use standard
- DESIRED = desired condition if flows are adequate to protect juvenile Lahontan cutthroat trout
- STD = Nevada State standard for Truckee River to protect beneficial uses

Table DSSAMt 6. Annual summary of simulated Truckee River quality compared to State of Nevada standards

Alternative: **No Action**

Year: **1992 CY**  
Run: 1992 (dry) - RT92N1: No Action  
Date of Run: 24-Feb-2004

Constituent	Statistic	Reach					
		Glend-McCar	McCar-Lockw	Lockw-Derby	Derby-Wadsw	Wadsw-Numan	Numan-Pyramid
pH	no. of days > STD	0	0	128	63	181	0
pH	no. of days < STD	0	0	0	0	0	0
Dissolved Oxygen	no. of days < STD (5 or 6 mg/L)	0	51	42	95	180	147
Dissolved Oxygen	no. of days < 5 mg/L*	0	51	40	59	147	127
Chloride	no. of days > STD#	366	124	124	267	271	340
Total Phosphorus	annual average	0.029	0.086	0.125	0.118	0.089	0.082
Ortho Phosphorus	no. of days > STD	0					
Total Nitrogen	annual average	0.332	0.630	0.777	0.693	0.598	0.594
Total Nitrogen	no. of days > STD	123	63	72	43	29	31
Nitrate Nitrogen	no. of days > STD	0	0	0	0	0	0
Nitrite Nitrogen	no. of days > STD	0	1	1	0	5	7
Ammonia Nitrogen (unionized)	no. of days > STD	0	0	0	0	0	0
Total Dissolved Solids	annual average#	142	195	228	235	433	594
Total Dissolved Solids	no. of days > STD#	337	124	121	62	300	355
Temperature	no. of days > STD fish flow not adequate	201	149	120	61	85	84
Temperature	no. of days > STD fish flow adequate ^	na	na	na	93	103	104
Temperature	no. of days > DESIRED fish flow not adequate	213	181	157	134	150	150
Temperature	no. of days > DESIRED fish flow adequate	213	181	157	172	186	186

Notes:

- no. of days = number of days of the year when the standard was exceeded within any hour of the day
- ^ = determined for period November through June = total of 273 days if flows are adequate for spawning
- na = not applicable
- \* = not a State Standard, included for information purposes only
- # = RMHQ (Recommended to Maintain High Water Quality) used in place of beneficial use standard
- DESIRED = desired condition if flows are adequate to protect juvenile Lahontan cutthroat trout
- STD = Nevada State standard for Truckee River to protect beneficial uses

Table DSSAMt 7. Annual summary of simulated Truckee River quality compared to State of Nevada standards

Alternative: **LWSA**
 Year: **1986 CY**  
 Run: **1986 (wet) - RT86L1: LWSA**  
 Date of Run: **24-Feb-2004**

Constituent	Statistic	Reach					
		Glend-McCar	McCar-Lockw	Lockw-Derby	Derby-Wadsw	Wadsw-Numan	Numan-Pyramid
pH	no. of days > STD	0	0	0	15	28	97
pH	no. of days < STD	0	0	0	0	0	0
Dissolved Oxygen	no. of days < STD (5 or 6 mg/L)	0	0	0	0	0	4
Dissolved Oxygen	no. of days < 5 mg/L*	0	0	0	0	0	4
Chloride	no. of days > STD#	63	0	0	0	0	0
Total Phosphorus	annual average	0.013	0.030	0.042	0.039	0.039	0.039
Ortho Phosphorus	no. of days > STD	0					
Total Nitrogen	annual average	0.179	0.273	0.335	0.319	0.324	0.329
	no. of days > STD	1	1	1	1	3	3
Nitrate Nitrogen	no. of days > STD	0	0	0	0	0	0
Nitrite Nitrogen	no. of days > STD	0	1	1	0	0	1
Ammonia Nitrogen (unionized)	no. of days > STD	0	0	0	0	1	1
Total Dissolved Solids	annual average#	69	92	108	106	119	126
	no. of days > STD#	2	0	0	0	0	0
Temperature	no. of days > STD fish flow not adequate	146	25	32	20	27	37
	no. of days > STD fish flow adequate ^	na	na	na	36	37	39
	no. of days > DESIRED fish flow not adequate	156	53	58	64	66	72
	no. of days > DESIRED fish flow adequate	156	53	58	100	103	106

Notes:

- no. of days = number of days of the year when the standard was exceeded within any hour of the day  
 ^ = determined for period November through June = total of 273 days if flows are adequate for spawning  
 na = not applicable  
 \* = not a State Standard, included for information purposes only  
 # = RMHQ (Recommended to Maintain High Water Quality) used in place of beneficial use standard  
 DESIRED = desired condition if flows are adequate to protect juvenile Lahontan cutthroat trout  
 STD = Nevada State standard for Truckee River to protect beneficial uses

Table DSSAMt 8. Annual summary of simulated Truckee River quality compared to State of Nevada standards

Alternative: **LWSA**Year: **1989 CY**  
Run: 1989 (median) - RT89L1: LWSA  
Date of Run: 29-Feb-2004

Constituent	Statistic	Reach					
		Glend-McCar	McCar-Lockw	Lockw-Derby	Derby-Wadsw	Wadsw-Numan	Numan-Pyramid
pH	no. of days > STD	0	0	26	86	147	136
pH	no. of days < STD	0	0	0	0	0	0
Dissolved Oxygen	no. of days < STD (5 or 6 mg/L)	0	0	0	4	10	26
Dissolved Oxygen	no. of days < 5 mg/L*	0	0	0	0	9	30
Chloride	no. of days > STD#	91	0	0	1	0	10
Total Phosphorus	annual average	0.013	0.013	0.014	0.043	0.055	0.050
Ortho Phosphorus	no. of days > STD	0					
Total Nitrogen	annual average	0.218	0.222	0.227	0.394	0.459	0.440
	no. of days > STD	1	1	1	2	3	1
Nitrate Nitrogen	no. of days > STD	0	0	0	0	0	0
Nitrite Nitrogen	no. of days > STD	0	1	1	0	0	0
Ammonia Nitrogen (unionized)	no. of days > STD	0	0	0	0	0	0
Total Dissolved Solids	annual average#	98	98	97	126	143	153
	no. of days > STD#	60	0	0	0	0	8
Temperature	no. of days > STD fish flow not adequate	154	13	27	21	51	62
	no. of days > STD fish flow adequate ^	na	na	na	77	80	82
	no. of days > DESIRED fish flow not adequate	160	51	61	86	98	107
	no. of days > DESIRED fish flow adequate	160	51	61	146	155	160

## Notes:

- no. of days = number of days of the year when the standard was exceeded within any hour of the day  
 ^ = determined for period November through June = total of 273 days if flows are adequate for spawning  
 na = not applicable  
 \* = not a State Standard, included for information purposes only  
 # = RMHQ (Recommended to Maintain High Water Quality) used in place of beneficial use standard  
 DESIRED = desired condition if flows are adequate to protect juvenile Lahontan cutthroat trout  
 STD = Nevada State standard for Truckee River to protect beneficial uses



Table DSSAMt 9. Annual summary of simulated Truckee River quality compared to State of Nevada standards

Alternative: **LWSA**

Year: **1992 CY (dry)**  
Run: **1992 (dry) - RT92L1: LWSA**  
Date of Run: **24-Feb-2004**

Constituent	Statistic	Reach					
		Glend-McCar	McCar-Lockw	Lockw-Derby	Derby-Wadsw	Wadsw-Numan	Numan-Pyramid
pH	no. of days > STD	0	0	103	54	182	0
pH	no. of days < STD	0	0	0	0	0	0
Dissolved Oxygen	no. of days < STD (5 or 6 mg/L)	0	46	39	97	177	149
Dissolved Oxygen	no. of days < 5 mg/L*	0	46	39	57	140	125
Chloride	no. of days > STD#	366	184	185	293	272	341
Total Phosphorus	annual average	0.029	0.085	0.123	0.115	0.088	0.081
Ortho Phosphorus	no. of days > STD	0					
Total Nitrogen	annual average	0.333	0.660	0.830	0.733	0.615	0.604
	no. of days > STD	123	93	93	65	30	32
Nitrate Nitrogen	no. of days > STD	0	0	0	0	0	0
Nitrite Nitrogen	no. of days > STD	0	1	1	0	5	7
Ammonia Nitrogen (unionized)	no. of days > STD	0	0	0	0	0	0
Total Dissolved Solids	annual average#	142	204	245	251	441	598
	no. of days > STD#	337	185	186	94	310	366
Temperature	no. of days > STD fish flow not adequate	201	149	119	61	85	84
	no. of days > STD fish flow adequate ^	na	na	na	93	103	104
	no. of days > DESIRED fish flow not adequate	213	181	156	134	150	150
	no. of days > DESIRED fish flow adequate	213	181	156	172	185	186

Notes:

- no. of days = number of days of the year when the standard was exceeded within any hour of the day
- ^ = determined for period November through June = total of 273 days if flows are adequate for spawning
- na = not applicable
- \* = not a State Standard, included for information purposes only
- # = RMHQ (Recommended to Maintain High Water Quality) used in place of beneficial use standard
- DESIRED = desired condition if flows are adequate to protect juvenile Lahontan cutthroat trout
- STD = Nevada State standard for Truckee River to protect beneficial uses

Table DSSAMt 10. Annual summary of simulated Truckee River quality compared to State of Nevada standards

Alternative: **TROA**
 Year: **1986 CY**  
 Run: **1986 (wet) - RT86T1: TROA**  
 Date of Run: **24-Feb-2004**

Constituent	Statistic	Reach					
		Glend-McCar	McCar-Lockw	Lockw-Derby	Derby-Wadsw	Wadsw-Numan	Numan-Pyramid
pH	no. of days > STD	0	0	0	13	17	102
pH	no. of days < STD	0	0	0	0	0	0
Dissolved Oxygen	no. of days < STD (5 or 6 mg/L)	0	0	0	0	0	0
Dissolved Oxygen	no. of days < 5 mg/L*	0	0	0	0	0	0
Chloride	no. of days > STD#	93	31	31	30	0	0
Total Phosphorus	annual average	0.014	0.032	0.046	0.043	0.043	0.042
Ortho Phosphorus	no. of days > STD	0					
Total Nitrogen	annual average	0.192	0.296	0.363	0.355	0.356	0.356
	no. of days > STD	1	1	1	1	3	4
Nitrate Nitrogen	no. of days > STD	0	0	0	0	0	0
Nitrite Nitrogen	no. of days > STD	0	1	1	0	0	0
Ammonia Nitrogen (unionized)	no. of days > STD	0	0	0	0	0	1
Total Dissolved Solids	annual average#	70	95	113	114	126	134
	no. of days > STD#	1	0	0	0	0	0
Temperature	no. of days > STD fish flow not adequate	131	21	29	13	22	30
	no. of days > STD fish flow adequate ^	na	na	na	35	36	38
	no. of days > DESIRED fish flow not adequate	137	47	55	60	65	66
	no. of days > DESIRED fish flow adequate	137	47	55	95	101	104

Notes:

- no. of days = number of days of the year when the standard was exceeded within any hour of the day  
 ^ = determined for period November through June = total of 273 days if flows are adequate for spawning  
 na = not applicable  
 \* = not a State Standard, included for information purposes only  
 # = RMHQ (Recommended to Maintain High Water Quality) used in place of beneficial use standard  
 DESIRED = desired condition if flows are adequate to protect juvenile Lahontan cutthroat trout  
 STD = Nevada State standard for Truckee River to protect beneficial uses

Table DSSAMt 11. Annual summary of simulated Truckee River quality compared to State of Nevada standards

Alternative: **TROA**

Year: **1989 CY**  
 Run: **1989 (median) - RT89T1: TROA**  
 Date of Run: **26-Feb-2004**

Constituent	Statistic	Reach					
		Glend-McCar	McCar-Lockw	Lockw-Derby	Derby-Wadsw	Wadsw-Numan	Numan-Pyramid
pH	no. of days > STD	0	0	19	81	140	141
pH	no. of days < STD	0	0	0	0	0	0
Dissolved Oxygen	no. of days < STD (5 or 6 mg/L)	0	0	0	4	10	15
Dissolved Oxygen	no. of days < 5 mg/L*	0	0	0	0	9	19
Chloride	no. of days > STD#	91	0	0	1	0	10
Total Phosphorus	annual average	0.016	0.039	0.055	0.049	0.047	0.046
Ortho Phosphorus	no. of days > STD	0					
Total Nitrogen	annual average	0.241	0.374	0.453	0.432	0.429	0.429
	no. of days > STD	1	1	1	2	3	1
Nitrate Nitrogen	no. of days > STD	0	0	0	0	0	0
Nitrite Nitrogen	no. of days > STD	0	1	1	0	0	0
Ammonia Nitrogen (unionized)	no. of days > STD	0	0	0	0	0	0
Total Dissolved Solids	annual average#	96	121	140	143	193	223
	no. of days > STD#	60	0	0	0	0	8
Temperature	no. of days > STD fish flow not adequate	154	14	28	21	53	62
	no. of days > STD fish flow adequate ^	na	na	na	77	79	82
	no. of days > DESIRED fish flow not adequate	157	51	63	89	99	107
	no. of days > DESIRED fish flow adequate	157	51	63	149	155	160

Notes:

no. of days = number of days of the year when the standard was exceeded within any hour of the day  
 ^ = determined for period November through June = total of 273 days if flows are adequate for spawning  
 na = not applicable  
 \* = not a State Standard, included for information purposes only  
 # = RMHQ (Recommended to Maintain High Water Quality) used in place of beneficial use standard  
 DESIRED = desired condition if flows are adequate to protect juvenile Lahontan cutthroat trout  
 STD = Nevada State standard for Truckee River to protect beneficial uses

Table DSSAMt 12. Annual summary of simulated Truckee River quality compared to State of Nevada standards

Alternative: **TROA**

Year: **1992 CY**  
Run: 1992 (dry) - RT92T1: TROA  
Date of Run: 24-Feb-2004

Constituent	Statistic	Reach					
		Glend-McCar	McCar-Lockw	Lockw-Derby	Derby-Wadsw	Wadsw-Numan	Numan-Pyramid
pH	no. of days > STD	0	0	99	49	143	24
pH	no. of days < STD	0	0	0	0	0	0
Dissolved Oxygen	no. of days < STD (5 or 6 mg/L)	0	1	3	65	126	148
Dissolved Oxygen	no. of days < 5 mg/L*	0	0	0	34	75	114
Chloride	no. of days > STD#	366	93	93	157	163	265
Total Phosphorus	annual average	0.027	0.078	0.114	0.108	0.087	0.083
Ortho Phosphorus	no. of days > STD	0					
Total Nitrogen	annual average	0.313	0.592	0.736	0.662	0.583	0.573
	no. of days > STD	123	63	64	40	9	9
Nitrate Nitrogen	no. of days > STD	0	0	0	0	0	0
Nitrite Nitrogen	no. of days > STD	0	1	1	0	0	0
Ammonia Nitrogen (unionized)	no. of days > STD	0	0	0	0	0	0
Total Dissolved Solids	annual average#	136	183	213	220	371	462
	no. of days > STD#	337	93	93	56	194	275
Temperature	no. of days > STD fish flow not adequate	189	81	87	49	75	78
	no. of days > STD fish flow adequate ^	na	na	na	90	98	100
	no. of days > DESIRED fish flow not adequate	191	139	133	125	145	146
	no. of days > DESIRED fish flow adequate	191	139	133	164	180	182

Notes:

- no. of days = number of days of the year when the standard was exceeded within any hour of the day
- ^ = determined for period November through June = total of 273 days if flows are adequate for spawning
- na = not applicable
- \* = not a State Standard, included for information purposes only
- # = RMHQ (Recommended to Maintain High Water Quality) used in place of beneficial use standard
- DESIRED = desired condition if flows are adequate to protect juvenile Lahontan cutthroat trout
- STD = Nevada State standard for Truckee River to protect beneficial uses

that population is expected to increase in the study area, which would increase municipal and industrial water demand. No Action serves as a baseline for comparison to the TROA Alternative.

Simulation of future conditions without TROA indicated warm temperatures for fish and water quality problems will exist minimally in wet and mostly in median and dry hydrologic conditions (Brock and Caupp, RCR04-5.0, 2004 and RCR04-6.0, 2004 ). Temperature, pH, DO, chloride, TDS, and total nitrogen violations will occur downstream from Reno. No Action is slightly worse than the current conditions due to larger future water demands, mostly due to population growth. The most severe conditions occur during dry hydrologic conditions. Without changes, current system operations will not adequately accommodate water quality in the future.

Annual summaries of major model water quality parameters under No Action are given in tables DSSAMt 4, 5, and 6 for each modeled reach downstream from Reno for wet (1986), median (1989), and dry (1992) calendar year conditions.

Annual summaries indicated that Truckee River TDS and chloride concentration standards downstream from Reno to Pyramid Lake, under dry year No Action conditions, are exceeded most of a dry year. Annual summaries indicated that total nitrogen in the reach from Lockwood to Derby Diversion Dam is exceeded about one quarter of the year.

## **X. LOCAL WATER SUPPLY ALTERNATIVE TEMPERATURE AND WATER QUALITY SIMULATIONS**

The Local Water Supply Alternative represents projected future conditions in the year 2033 without a Truckee River operating agreement; however, with some likely changes in system operation. Therefore, LWSA and No Action do not vary greatly. The primary differences between LWSA and No Action would be the source of water used for municipal and industrial purposes, extent of water conservation, implementation of a groundwater recharge program in Truckee Meadows, and assumptions regarding governmental decisions concerning approval of new water supply proposals.

Simulation of future conditions under LWSA indicated warm temperatures for fish and water quality problems will exist primarily in wet, average, and dry conditions (Brock and Caupp, RCR04-9.0, 2004 and RCR04-10.0, 2004). Temperature, pH, DO, chloride, TDS, and total nitrogen violations will occur downstream from Reno. LWSA is slightly worse than the current conditions due to larger future water demands mostly due to population growth. The most severe conditions occur during dry conditions.

Annual summaries of major model water quality parameters under the LWSA are given in tables 7, 8, and 9 for each modeled reach downstream from Reno for wet (1986), median (1989), and dry (1992) calendar year conditions.

Under dry year LWSA conditions, annual summaries indicated that TDS and chloride concentration standards downstream from Reno are exceeded more often than under



No Action. Under dry year LWSA conditions, annual summaries indicated that total nitrogen standards in the reach from Lockwood to Derby Diversion Dam are exceeded about one quarter of the year and again are exceeded more often than under No Action.

## **XI. TROA ALTERNATIVE TEMPERATURE AND WATER QUALITY SIMULATIONS**

The TROA Alternative represents a blended Truckee River operating agreement to accommodate the concerns of several negotiating parties. This alternative concentrates on implementing the requirements of the Preliminary Settlement Act (PSA). It provides drought relief for the Reno/Sparks area and enhances spawning flows for threatened and endangered fishes of Pyramid Lake. Under this alternative, current mandatory minimum instream flows would be met.

TROA provides a compromise of several resources and accommodates the needs of many users. Therefore, river temperatures and water quality would not be expected to be optimum for the nonhuman environment under TROA. However, from a water quality perspective, modeling indicated that TROA is better than No Action in most instances (Brock and Caupp, RCR04-7.0, 2004 and RCR04-8.0, 2004). The TROA Alternative appears to be the optimum alternative for the human environment.

Annual summaries of major model water quality parameters under TROA are given in tables DSSAMt 10, 11, and 12 for each modeled reach downstream from Reno for wet (1986), median (1989), and dry (1992) calendar year conditions.

Under dry year TROA conditions, annual summaries indicated that TDS and chloride concentration standards are exceeded less often than under No Action. Annual summaries indicated that total nitrogen standards in the reach from Lockwood to Derby Diversion Dam are exceeded less than one third of the year but less often than under No Action.

## **XII. SIMULATED VERSUS PREFERRED WATER TEMPERATURES FOR FISH**

Tables and plots summarizing fish temperature preferences for appropriate life stages of Truckee River fishes were prepared for the current condition simulation (Brock and Caupp, RCR04-3.0, 2004 ) and for the alternatives (Brock and Caupp, RCR04-5.0, 2004 , RCR04-7.0, 2004, and RCR04\_9.0, 2004). These included temperature preference windows on plots of daily temperature for facilitating interpretation of the wealth of fish life stage information. Simulated daily instantaneous maximum temperatures indicated acute temperature effects on fish, whereas 7-day moving averages of maximum, mean, and minimum temperatures indicated chronic temperature effects. Life stages were separated into adult migration, spawning, incubation, larvae or rearing, juvenile maintenance, and adult maintenance. Fish species included rainbow trout, cui-ui, Lahontan cutthroat trout, mountain whitefish, and brown trout. Summary tables of the information for extremely wet, median,

and extremely dry ending years of a 5-year period for the current conditions and the alternatives (12 tables of five fishes and associated life stages from Chapter 6 of the Brock and Caupp fish temperature books) are included in the following WQ Appendix DSSAMt tables 13 through DSSAMt 24.

Simulated river temperatures were compared to preferred ranges. For each species and life stage of fish, the number of days that the simulated temperatures were within the recommended range was counted. The temperature was counted as having met the preferred criteria if the temperature was less than or equal to the maximum preferred temperature and greater than or equal to the minimum preferred temperature. These criteria are exceeded if the temperature curves pass through the bottom or top of the thermal box. The number of days that recommended temperatures were met during each fish life stage was given for the daily instantaneous maximum and maximum, mean, and minimum 7-day moving averages at six locations: East McCarron, Lockwood, Clark, Painted Rock, Dead Ox, and Marble Bluff Dam. These locations correspond to locations listed in the Nevada water temperature standards or critical locations determined by fishery biologists.

In general, during average and dry current and future conditions, recommended temperatures are exceeded for cold water and cool water fish. Recommended temperatures for warm water species such as cui-ui are exceeded, but much less frequently. During wet years in which cui-ui spawning runs occur, temperature requirements are often met during the high spring spawning flows. However, once flows are greatly reduced during summer and fall, preferred temperatures for cold water fish such as trout are exceeded frequently. Differences between alternatives were minimal in comparison to differences between wet, median, and dry years.

Summary tables of the water quality books are shown below as Water Quality Appendix DSSAMt tables 13 – 24.

Table DSSAMt 13. No. of days recommended temperatures are met for Truckee R. Fishes

**Truckee R. 1986wy (extremely wet) - Current Condition**

**Rainbow Trout**

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg			
		Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN
Adult Migration	46	39	41	41	33	35	38	44	37	31	36	46	39	31	36	43	40	29	30	38	41	25	28	36	40
Spawning	32	25	27	24	13	21	24	29	19	17	22	30	20	17	22	27	21	15	16	23	28	11	14	22	27
Incubation	62	32	36	40	39	24	27	44	37	20	22	40	37	20	22	36	38	15	16	23	38	11	14	22	37
Rearing	183	178	183	176	173	181	183	183	183	177	183	183	178	178	183	183	177	162	167	183	176	146	158	183	175
Juvenile Maint.	365	355	360	342	335	363	365	365	365	359	365	366	349	360	365	361	347	344	349	358	345	328	340	358	342
Adult Maint.	365	355	360	342	335	363	365	365	365	359	365	365	349	360	365	361	347	344	349	358	345	328	340	358	342

**Cui-ui**

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg			
		Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN
Adult Migration	150	78	83	46	35	83	85	54	38	74	79	58	44	70	78	59	45	63	63	58	47	58	59	51	46
Spawning	91	64	69	46	35	68	70	52	38	60	65	56	44	57	65	57	45	52	52	56	47	48	49	49	45
Incubation	91	71	71	46	35	76	76	52	38	77	77	56	44	77	77	57	45	74	76	62	47	74	75	65	47
Larvae	62	55	56	47	36	56	57	51	39	57	58	52	45	57	58	52	46	54	57	55	48	53	54	55	48

**Lahontan Cutthroat Trout**

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg			
		Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN
Adult Migration	136	100	102	114	102	98	102	107	111	96	101	98	106	95	100	96	101	94	98	94	95	92	99	93	92
Spawning	91	59	60	59	44	56	57	60	48	53	57	51	50	55	58	50	50	53	56	48	41	52	57	49	41
Incubation	91	59	60	59	44	56	57	60	48	53	57	51	50	55	58	50	50	53	56	48	41	52	57	49	41
Juvenile Maint.	107	88	95	96	92	94	98	103	96	81	91	105	96	82	92	107	97	74	83	107	98	57	60	107	96
Adult Maint.	365	338	338	331	309	349	349	339	334	340	343	336	333	338	339	335	333	326	330	335	332	323	332	335	331

**Mountain Whitefish**

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg			
		Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN
Adult Migration	62	37	39	48	51	31	38	42	48	37	36	42	47	41	36	41	47	32	32	42	48	20	16	41	47
Spawning	47	24	23	36	33	28	26	30	39	32	28	27	37	27	24	27	35	27	21	29	35	26	22	28	34
Incubation	78	37	36	47	42	55	53	45	52	54	50	40	49	45	42	40	47	47	41	42	47	53	49	41	46
Incubation	74	74	74	74	70	71	74	74	74	72	74	74	74	74	74	74	74	74	74	74	74	74	74	74	74
Rearing	154	135	136	154	154	135	136	154	154	130	131	151	154	130	131	151	154	128	126	149	154	122	123	148	154
Juvenile Maint.	365	309	307	342	335	316	314	360	365	307	302	342	349	308	302	335	347	301	295	330	345	295	293	327	342
Adult Maint.	365	309	307	342	335	316	314	360	365	307	302	342	349	308	302	335	347	301	295	330	345	295	293	327	342

**Brown Trout**

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg			
		Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN
Adult Migration	62	37	39	48	51	31	38	42	48	37	36	42	47	41	36	41	47	32	32	42	48	20	16	41	47
Spawning	32	15	14	30	29	13	11	20	31	17	13	18	31	17	14	20	30	16	10	24	31	11	7	22	31
Incubation	78	37	36	47	42	55	53	45	52	54	50	40	49	45	42	40	47	47	41	42	47	53	49	41	46
Incubation	74	74	74	74	70	71	74	74	74	72	74	74	74	74	74	74	74	74	74	74	74	74	74	74	74
Rearing	154	151	154	154	154	153	154	154	154	150	154	154	154	151	154	154	154	147	154	154	154	139	149	154	154
Juvenile Maint.	365	355	360	342	335	363	365	365	365	359	365	365	349	360	365	361	347	344	349	358	345	328	340	358	342
Adult Maint.	365	355	360	342	335	363	365	365	365	359	365	365	349	360	365	361	347	344	349	358	345	328	340	358	342

Number of Annual Degree Days (deg C)

3,375

3,677

3,757

3,772

3,838

3,894

RCR: F:\TROA\_03\Fish\Fishsum\FishTempSum4.wb3

rev 18-Feb-2004

Date of Truckee River Operating Model Analysis:

10-Feb-04

RCR: 1

\*\*Provisional Data\*\*

17-Feb-2004

Table DSSAMt 14. No. of days recommended temperatures are met for Truckee R. Fishes

Truckee R. 1966wy (median) - Current Condition

Rainbow Trout

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg			
		MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN
Adult Migration	46	40	44	46	46	36	43	46	46	30	34	46	46	29	32	45	46	20	22	35	46	6	0	31	46
Spawning	32	26	30	32	11	22	29	32	21	17	20	32	32	17	18	31	32	11	11	21	32	3	0	17	32
Incubation	62	34	32	55	41	24	29	50	47	18	20	32	52	17	18	31	47	11	11	21	32	3	0	17	32
Rearing	183	182	183	183	183	183	183	183	183	182	183	183	183	181	183	183	183	162	163	183	183	141	143	183	183
Juvenile Maint.	365	364	365	365	354	365	365	365	365	364	365	365	365	363	365	365	365	344	345	365	365	323	325	365	361
Adult Maint.	365	364	365	365	354	365	365	365	365	364	365	365	365	363	365	365	365	344	345	365	365	323	325	365	361

Cui-ui

Life Stage	Total # Days	East McCarran 96.7 km				East McCarran 96.7 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg			
		MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN
Adult Migration	150	89	90	88	52	87	91	91	75	79	85	83	86	77	84	80	88	70	75	79	83	55	58	73	83
Spawning	91	76	77	86	52	72	76	88	75	68	74	80	85	67	74	77	87	57	61	75	81	41	41	69	81
Incubation	91	86	89	88	52	86	88	91	75	84	88	91	87	82	88	90	89	74	75	89	91	69	74	84	91
Larvae	62	57	59	62	51	57	58	62	62	55	58	62	62	53	58	60	62	44	45	59	62	39	44	54	62

Lahontan Cutthroat Trout

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg			
		MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN
Adult Migration	136	65	72	94	95	59	65	99	96	53	53	83	92	49	51	80	88	32	36	51	79	33	31	42	75
Spawning	91	48	57	77	74	41	44	75	85	36	31	64	76	35	30	63	75	15	17	48	65	10	5	32	63
Incubation	91	48	57	77	74	41	44	75	85	36	31	64	76	35	30	63	75	15	17	48	65	10	5	32	63
Juvenile Maint.	107	102	107	107	100	103	107	107	107	96	100	107	107	93	99	107	107	72	72	107	107	53	54	107	107
Adult Maint.	365	364	364	343	320	365	365	361	346	362	362	354	343	360	360	354	341	343	346	351	336	325	325	352	328

Mountain Whitefish

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg			
		MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN
Adult Migration	62	26	18	43	62	23	18	33	50	20	17	26	36	20	17	23	34	18	16	17	26	11	9	17	26
Spawning	47	32	28	47	47	29	26	45	47	33	28	44	46	33	29	42	45	33	25	36	44	29	24	35	44
Incubation	78	63	59	74	67	60	57	76	73	64	59	72	69	64	60	69	68	61	53	60	65	60	55	57	63
Incubation	74	71	74	64	52	68	71	74	69	71	74	74	68	72	74	74	68	71	74	74	68	71	74	74	68
Rearing	154	138	142	154	152	138	142	154	154	134	138	154	154	134	137	150	154	110	116	140	154	106	106	138	154
Juvenile Maint.	365	332	333	365	354	332	333	365	365	325	326	350	365	324	324	343	365	289	297	328	364	275	278	325	361
Adult Maint.	365	332	333	365	354	332	333	365	365	325	326	350	365	324	324	343	365	289	297	328	364	275	278	325	361

Brown Trout

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg			
		MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN
Adult Migration	62	26	18	43	62	23	18	33	50	20	17	26	36	20	17	23	34	18	16	17	26	11	9	17	26
Spawning	32	17	13	32	32	14	11	30	32	18	13	29	31	18	14	27	30	18	10	21	29	14	9	20	29
Incubation	78	63	59	74	67	60	57	76	73	64	59	72	69	64	60	69	68	61	53	60	65	60	55	57	63
Incubation	74	71	74	64	52	68	71	74	69	71	74	74	68	72	74	74	68	71	74	74	68	71	74	74	68
Rearing	154	153	154	154	152	154	154	154	154	154	154	154	154	154	154	154	154	148	148	154	154	133	137	154	154
Juvenile Maint.	365	364	365	365	354	365	365	365	365	364	365	365	365	363	365	365	365	344	345	365	365	323	325	365	361
Adult Maint.	365	364	365	365	354	365	365	365	365	364	365	365	365	363	365	365	365	344	345	365	365	323	325	365	361

Number of Annual Degree Days (deg C)

3,485

3,812

3,989

4,031

4,244

4,323

RCR | J:\TRQA\_03\Fish-Fishsum\FishTempSum4.tbl

rev 18-Feb-2004

Date of Truckee River Operating Model Analysis:

10-Feb-04

RCR: 1

\*\*Provisional Data\*\*

17-Feb-2004

Table DSSAMt 15. No. of days recommended temperatures are met for Truckee R. Fishes

## Truckee R. 1992wy (extremely dry) - Current Condition

## Rainbow Trout

Life Stage	Total # Days	East McCarran 96.7 km				Wood 106				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg			
		Inst.	MAX	MEAN	MIN	Inst.	MAX	MEAN	MIN	Inst.	MAX	MEAN	MIN	Inst.	MAX	MEAN	MIN	Inst.	MAX	MEAN	MIN	Inst.	MAX	MEAN	MIN
Adult Migration	46	3	0	14	34	1	0	5	26	1	0	3	17	0	0	3	17	1	0	3	13	0	0	3	25
Spawning	32	2	0	8	20	0	0	0	12	0	0	0	10	0	0	0	10	1	0	0	6	0	0	0	13
Incubation	62	2	0	13	44	0	0	0	15	0	0	0	18	0	0	0	16	1	0	0	12	0	0	0	27
Rearing	183	144	145	183	183	123	128	183	183	132	137	183	183	138	140	183	183	154	155	183	183	129	133	183	183
Juvenile Maint	365	326	327	365	348	305	310	365	365	314	319	365	365	320	322	365	365	335	337	365	362	306	315	365	365
Adult Maint	365	326	327	365	348	305	310	365	365	314	319	365	365	320	322	365	365	335	337	365	362	306	315	365	365

## Cui-ui

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg			
		Inst.	MAX	MEAN	MIN	Inst.	MAX	MEAN	MIN	Inst.	MAX	MEAN	MIN	Inst.	MAX	MEAN	MIN	Inst.	MAX	MEAN	MIN	Inst.	MAX	MEAN	MIN
Adult Migration	150	62	61	67	67	66	68	77	87	64	66	63	68	63	65	57	68	63	65	55	66	53	60	68	74
Spawning	91	30	29	64	67	27	29	48	64	29	31	42	65	28	30	39	64	29	31	39	61	19	24	42	69
Incubation	91	58	61	87	71	54	55	70	86	52	55	73	91	47	49	72	91	49	49	68	91	38	36	72	91
Larvae	62	29	31	58	62	24	25	40	57	22	25	44	62	18	19	43	62	20	19	39	62	9	6	43	62

## Lahontan Cutthroat Trout

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg			
		Inst.	MAX	MEAN	MIN	Inst.	MAX	MEAN	MIN	Inst.	MAX	MEAN	MIN	Inst.	MAX	MEAN	MIN	Inst.	MAX	MEAN	MIN	Inst.	MAX	MEAN	MIN
Adult Migration	136	54	67	67	87	52	59	72	77	56	60	70	67	49	52	66	68	50	54	65	67	37	35	68	71
Spawning	91	17	24	31	65	17	24	28	36	19	23	28	34	14	11	28	35	14	14	29	34	8	7	28	42
Incubation	91	17	24	31	65	17	24	28	36	19	23	28	34	14	11	28	35	14	14	29	34	8	7	28	42
Juvenile Maint	107	64	63	107	107	32	36	90	107	56	59	97	107	57	60	96	107	67	68	105	107	52	54	107	107
Adult Maint	365	328	337	324	309	311	317	365	365	323	323	365	357	329	328	351	329	341	343	342	315	311	318	361	337

## Mountain Whitefish

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg			
		Inst.	MAX	MEAN	MIN	Inst.	MAX	MEAN	MIN	Inst.	MAX	MEAN	MIN	Inst.	MAX	MEAN	MIN	Inst.	MAX	MEAN	MIN	Inst.	MAX	MEAN	MIN
Adult Migration	62	11	8	11	43	3	0	4	6	9	6	8	9	9	6	8	10	10	7	8	10	9	6	8	11
Spawning	47	31	27	35	39	18	15	16	24	27	23	28	32	28	24	30	36	30	28	35	36	25	21	31	36
Incubation	78	62	58	58	45	49	46	47	55	58	54	59	63	59	55	61	61	61	59	64	58	56	52	62	64
Incubation	74	59	64	63	52	55	60	73	74	60	62	74	74	58	58	73	64	61	63	64	55	54	55	71	64
Rearing	154	101	101	151	154	92	94	115	151	98	96	124	152	94	93	124	151	95	97	127	152	80	78	132	154
Juvenile Maint	365	288	286	359	348	229	226	299	348	251	242	312	355	253	248	312	351	275	279	316	351	231	221	321	365
Adult Maint	365	288	286	359	348	229	226	299	348	251	242	312	355	253	248	312	351	275	279	316	351	231	221	321	365

## Brown Trout

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg			
		Inst.	MAX	MEAN	MIN	Inst.	MAX	MEAN	MIN	Inst.	MAX	MEAN	MIN	Inst.	MAX	MEAN	MIN	Inst.	MAX	MEAN	MIN	Inst.	MAX	MEAN	MIN
Adult Migration	62	11	8	11	43	3	0	4	6	9	6	8	9	9	6	8	10	10	7	8	10	9	6	8	11
Spawning	32	16	12	20	24	3	0	1	9	12	8	13	17	13	9	15	21	15	13	20	21	10	7	16	21
Incubation	78	62	58	58	45	49	46	47	55	58	54	59	63	59	55	61	61	61	59	64	58	56	52	62	64
Incubation	74	59	64	63	52	55	60	73	74	60	62	74	74	58	58	73	64	61	63	64	55	54	55	71	64
Rearing	154	128	134	154	154	117	122	154	154	121	130	154	154	125	133	154	154	137	143	154	154	114	126	154	154
Juvenile Maint	365	326	327	365	348	305	310	365	365	314	319	365	365	320	322	365	365	335	337	365	362	306	315	365	365
Adult Maint	365	326	327	365	348	305	310	365	365	314	319	365	365	320	322	365	365	335	337	365	362	306	315	365	365

Number of Annual Degree Days (deg C)

4,127

5,197

4,812

4,722

4,816

4,715

RCR: 1:TRDA\_03-Fish/Fishsum/FishTempSum4.m3

rev: 18-Feb-2004

Date of Truckee River Operating Model Analysis:

10-Feb-04

RCR: 1

\*\*Provisional Data\*\*

17-Feb-2004



Table DSSAMt 16. No. of days recommended temperatures are met for Truckee R. Fishes

## Truckee R. 1986wy (extremely wet) - No Action

### Rainbow Trout

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg			
		MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN
Adult Migration	46	39	41	41	33	32	38	46	38	31	36	45	40	31	35	43	40	28	30	38	41	25	28	36	39
Spawning	32	25	27	24	13	18	24	29	19	17	22	29	20	17	21	27	23	14	16	24	28	11	14	22	27
Incubation	62	32	36	40	39	21	27	44	37	19	22	39	37	20	21	36	40	14	16	24	38	11	14	22	37
Rearing	183	178	183	176	173	181	183	183	183	174	177	183	183	175	180	183	183	162	167	183	182	146	158	183	178
Juvenile Maint.	365	356	361	342	335	363	365	365	365	356	359	365	365	357	362	365	365	344	349	365	359	328	340	365	351
Adult Maint.	365	356	361	342	335	363	365	365	365	356	359	365	365	357	362	365	365	344	349	365	359	328	340	365	351

### Cui-ui

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg			
		MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN
Adult Migration	150	78	83	48	35	86	88	56	41	73	78	62	45	72	77	62	45	64	64	58	49	61	62	52	46
Spawning	91	64	69	46	35	70	72	54	41	59	64	60	45	58	63	60	45	53	53	56	49	50	51	48	46
Incubation	91	71	71	46	35	78	78	54	41	78	78	60	45	78	78	61	45	75	77	64	49	76	77	65	48
Larvae	62	55	56	47	36	57	58	52	42	57	58	54	46	57	58	54	46	55	58	55	49	54	55	55	46

### Lahontan Cutthroat Trout

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg			
		MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN
Adult Migration	136	100	102	114	102	98	102	109	113	96	102	98	107	96	101	96	103	94	100	94	96	91	99	94	94
Spawning	91	59	60	59	44	56	57	61	48	53	57	51	51	54	57	51	50	55	57	49	44	51	57	52	43
Incubation	91	59	60	59	44	56	57	61	48	53	57	51	51	54	57	51	50	55	57	49	44	51	57	52	43
Juvenile Maint.	107	89	95	96	92	92	98	104	96	78	90	107	97	81	91	107	97	75	83	107	98	57	60	107	96
Adult Maint.	365	339	339	332	315	365	365	362	343	358	362	345	336	354	366	342	336	336	339	338	334	330	340	337	333

### Mountain Whitefish

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg			
		MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN
Adult Migration	62	38	39	48	51	30	36	41	48	33	36	41	47	40	36	41	46	29	32	41	46	21	18	43	47
Spawning	47	24	22	36	33	25	22	33	47	26	23	33	39	29	27	30	37	29	24	30	36	25	22	31	36
Incubation	78	37	35	47	42	56	53	64	78	57	54	59	55	58	56	53	50	56	51	49	49	55	52	50	48
Incubation	74	74	74	74	72	68	74	74	74	71	74	74	74	72	74	74	74	74	74	74	74	74	74	74	74
Rearing	154	135	136	154	154	135	136	154	154	130	129	151	154	130	130	151	154	126	125	149	154	122	123	148	154
Juvenile Maint.	365	310	308	342	335	315	314	355	365	306	300	340	365	307	301	339	365	301	295	336	359	295	293	334	351
Adult Maint.	365	310	308	342	335	315	314	355	365	306	300	340	365	307	301	339	365	301	295	336	359	295	293	334	351

### Brown Trout

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg			
		MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN
Adult Migration	62	38	39	48	51	30	36	41	48	33	36	41	47	40	36	41	46	29	32	41	46	21	18	43	47
Spawning	32	15	13	30	29	10	7	18	32	11	8	18	31	14	12	18	30	14	9	22	31	10	7	22	32
Incubation	78	37	35	47	42	56	53	64	78	57	54	59	55	58	56	53	50	56	51	49	49	55	52	50	48
Incubation	74	74	74	74	72	68	74	74	74	71	74	74	74	72	74	74	74	74	74	74	74	74	74	74	74
Rearing	154	151	154	154	154	153	154	154	154	149	154	154	154	149	154	154	154	147	154	154	154	139	149	154	154
Juvenile Maint.	365	356	361	342	335	363	365	365	365	356	359	365	365	357	362	365	365	344	349	365	359	328	340	365	351
Adult Maint.	365	356	361	342	335	363	365	365	365	356	359	365	365	357	362	365	365	344	349	365	359	328	340	365	351

Number of Annual Degree Days (deg C)

33

38

364

373

313

342

RCR: f:\TROA\_03\Fish\Fishsum\FishTempSum4 web3

rev 18 Feb 2004

Date of Truckee River Operating Model Analysis:

10-Feb-04

RCR: 1

\*\*Provisional Data\*\*

17-Feb-2004

Table DSSAMt 17. No. of days recommended temperatures are met for Truckee R. Fishes

## Truckee R. 1966wy (median) - No Action

## Rainbow Trout

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst.	MAX	7-day moving avg	MIN	Inst.	MAX	7-day moving avg	MIN	Inst.	MAX	7-day moving avg	MIN	Inst.	MAX	7-day moving avg	MIN	Inst.	MAX	7-day moving avg	MIN	Inst.	MAX	7-day moving avg	MIN
Adult Migration	46	36	41	46	46	30	35	46	46	21	28	45	46	21	28	40	45	7	1	31	43	2	0	27	42
Spawning	32	22	27	32	16	18	21	32	28	12	14	31	32	12	14	26	32	4	1	17	29	1	0	13	28
Incubation	62	24	27	55	46	19	21	44	54	12	14	31	47	12	14	26	37	4	1	17	29	1	0	13	28
Rearing	183	182	183	183	183	183	183	183	183	181	183	183	183	180	183	183	183	161	182	183	183	139	141	183	183
Juvenile Maint.	365	364	365	360	341	365	365	365	365	363	365	365	365	362	365	365	365	343	344	365	365	321	323	365	365
Adult Maint.	365	364	365	360	341	365	365	365	365	363	365	365	365	362	365	365	365	343	344	365	365	321	323	365	365

## Cui-ui

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst.	MAX	7-day moving avg	MIN	Inst.	MAX	7-day moving avg	MIN	Inst.	MAX	7-day moving avg	MIN	Inst.	MAX	7-day moving avg	MIN	Inst.	MAX	7-day moving avg	MIN	Inst.	MAX	7-day moving avg	MIN
Adult Migration	150	87	90	87	57	89	95	95	88	77	82	84	90	78	79	82	89	59	70	71	79	43	41	68	82
Spawning	91	73	76	85	57	70	75	87	85	62	66	77	88	61	64	76	87	46	55	67	77	29	23	63	80
Incubation	91	86	88	88	57	85	88	91	85	82	87	90	90	82	86	90	91	71	75	87	91	62	65	82	91
Larvae	62	57	58	62	55	56	58	62	62	53	57	60	62	52	56	60	62	41	45	57	62	32	35	52	62

## Lahontan Cutthroat Trout

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst.	MAX	7-day moving avg	MIN	Inst.	MAX	7-day moving avg	MIN	Inst.	MAX	7-day moving avg	MIN	Inst.	MAX	7-day moving avg	MIN	Inst.	MAX	7-day moving avg	MIN	Inst.	MAX	7-day moving avg	MIN
Adult Migration	136	58	63	94	96	59	60	97	102	44	50	73	90	42	48	65	84	37	35	42	70	35	34	41	67
Spawning	91	40	46	76	74	32	31	70	82	24	23	58	75	22	23	54	69	11	12	29	61	6	0	23	60
Incubation	91	40	46	76	74	32	31	70	82	24	23	58	75	22	23	54	69	11	12	29	61	6	0	23	60
Juvenile Maint.	107	102	107	107	101	103	107	107	107	94	100	107	107	94	97	107	107	71	71	107	107	47	50	107	107
Adult Maint.	365	356	356	327	301	365	365	364	352	365	365	365	342	364	364	352	339	346	349	347	332	324	328	344	324

## Mountain Whitefish

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst.	MAX	7-day moving avg	MIN	Inst.	MAX	7-day moving avg	MIN	Inst.	MAX	7-day moving avg	MIN	Inst.	MAX	7-day moving avg	MIN	Inst.	MAX	7-day moving avg	MIN	Inst.	MAX	7-day moving avg	MIN
Adult Migration	62	22	17	42	62	20	17	19	45	18	16	18	31	18	16	17	30	15	15	16	20	9	7	16	22
Spawning	47	29	31	46	47	24	18	45	46	31	27	39	45	31	28	38	45	31	24	34	44	28	22	33	44
Incubation	78	60	62	75	68	55	49	76	77	62	58	70	73	62	59	69	73	62	55	62	67	59	53	59	65
Incubation	74	64	68	53	28	65	70	74	74	71	73	74	67	71	74	73	66	71	74	71	63	71	73	70	58
Rearing	154	137	141	154	153	135	139	154	154	132	135	149	154	132	135	145	154	110	107	140	154	102	104	137	154
Juvenile Maint.	365	331	332	360	341	330	330	365	365	323	323	346	365	322	323	338	365	290	288	328	365	271	276	325	365
Adult Maint.	365	331	332	360	341	330	330	365	365	323	323	346	365	322	323	338	365	290	288	328	365	271	276	325	365

## Brown Trout

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst.	MAX	7-day moving avg	MIN	Inst.	MAX	7-day moving avg	MIN	Inst.	MAX	7-day moving avg	MIN	Inst.	MAX	7-day moving avg	MIN	Inst.	MAX	7-day moving avg	MIN	Inst.	MAX	7-day moving avg	MIN
Adult Migration	62	22	17	42	62	20	17	19	45	18	16	18	31	18	16	17	30	15	15	16	20	9	7	16	22
Spawning	32	14	16	31	32	9	5	30	31	16	12	24	30	16	13	23	30	16	9	19	29	13	7	18	29
Incubation	78	60	62	75	68	55	49	76	77	62	58	70	73	62	59	69	73	62	55	62	67	59	53	59	65
Incubation	74	64	68	53	28	65	70	74	74	71	73	74	67	71	74	73	66	71	74	71	63	71	73	70	58
Rearing	154	153	154	154	153	154	154	154	154	153	154	154	154	153	154	154	154	147	147	154	154	131	135	154	154
Juvenile Maint.	365	364	365	360	341	365	365	365	365	363	365	365	365	362	365	365	365	343	344	365	365	321	323	365	365
Adult Maint.	365	364	365	360	341	365	365	365	365	363	365	365	365	362	365	365	365	343	344	365	365	321	323	365	365

Number of Annual Degree Days (deg C)

366

363

4,18

4,173

4,44

4,413

RCR \TROA\_03\Fish\Fishsum\FishTempSum4.wb3

rev '8 Feb 2004

Date of Truckee River Operating Model Analysis:

10-Feb-04

RCR: 1

\*\*Provisional Data\*\*

17-Feb-2004

Table DSSAMt 18. No. of days recommended temperatures are met for Truckee R. Fishes

# Truckee R. 1992wy (extremely dry) - No Action

## Rainbow Trout

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg			
		MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN
Adult Migration	46	3	0	12	33	1	0	1	19	0	0	2	11	0	0	2	11	1	0	3	12	0	0	3	25
Spawning	32	2	0	6	19	0	0	0	11	0	0	0	4	0	0	0	4	1	0	0	5	0	0	0	13
Incubation	62	2	0	11	42	0	0	0	11	0	0	0	9	0	0	0	9	1	0	0	11	0	0	0	28
Rearing	183	143	148	183	183	126	129	183	183	133	136	183	183	142	141	183	183	153	154	183	183	129	131	183	183
Juvenile Maint.	365	325	330	365	350	308	311	365	365	315	318	365	365	324	323	365	365	334	336	365	365	306	313	365	365
Adult Maint.	365	325	330	365	350	308	311	365	365	315	318	365	365	324	323	365	365	334	336	365	365	306	313	365	365

## Cui-ui

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg			
		MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN
Adult Migration	150	61	60	66	69	82	84	83	95	65	66	71	78	63	65	67	66	63	65	69	67	53	60	68	73
Spawning	91	29	28	63	69	26	28	39	62	29	30	37	61	28	30	37	58	29	31	40	61	19	24	41	68
Incubation	91	57	61	67	74	49	50	69	81	48	51	71	91	50	52	69	88	50	51	69	91	38	36	71	91
Larvae	62	28	31	58	62	19	20	39	52	18	21	42	62	21	22	40	59	21	21	40	62	9	6	42	62

## Lahontan Cutthroat Trout

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg			
		MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN
Adult Migration	136	52	60	66	85	44	42	70	73	48	53	71	74	46	51	72	67	50	53	65	67	37	35	68	72
Spawning	91	14	24	31	65	9	12	26	32	15	15	28	31	14	14	28	32	14	13	29	34	8	7	28	41
Incubation	91	14	24	31	65	9	12	26	32	15	15	28	31	14	14	28	32	14	13	29	34	8	7	28	41
Juvenile Maint.	107	65	65	107	107	32	32	90	107	55	59	96	107	59	61	95	107	66	68	107	107	53	54	107	107
Adult Maint.	365	331	335	325	309	313	318	365	365	323	323	365	365	333	331	365	365	342	345	345	326	311	318	364	338

## Mountain Whitefish

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg			
		MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN
Adult Migration	62	11	8	11	42	1	0	3	5	9	6	7	9	9	6	7	9	10	7	8	10	9	6	8	11
Spawning	47	30	27	34	39	3	0	2	13	25	19	25	29	26	22	27	30	30	28	35	35	25	21	31	36
Incubation	78	61	58	58	45	15	12	33	44	56	50	56	60	57	53	58	61	61	59	66	60	56	52	62	64
Incubation	74	59	63	63	52	26	24	48	65	55	55	65	74	56	56	70	74	60	63	66	57	54	55	71	66
Rearing	154	102	102	150	154	90	91	115	152	94	95	123	152	96	98	123	149	94	98	127	152	80	78	132	154
Juvenile Maint.	365	288	286	354	350	226	223	290	348	244	239	311	354	256	254	311	348	273	280	316	354	231	221	323	365
Adult Maint.	365	288	286	354	350	226	223	290	348	244	239	311	354	256	254	311	348	273	280	316	354	231	221	323	365

## Brown Trout

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg			
		MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN
Adult Migration	62	11	8	11	42	1	0	3	5	9	6	7	9	9	6	7	9	10	7	8	10	9	6	8	11
Spawning	32	15	12	19	24	0	0	0	0	10	6	10	14	11	8	12	15	15	13	20	20	10	7	16	21
Incubation	78	61	58	58	45	15	12	33	44	56	50	56	60	57	53	58	61	61	59	66	60	56	52	62	64
Incubation	74	59	63	63	52	26	24	48	65	55	55	65	74	56	56	70	74	60	63	66	57	54	55	71	66
Rearing	154	128	140	154	154	119	123	154	154	122	129	154	154	129	134	154	154	137	143	154	154	114	124	154	154
Juvenile Maint.	365	325	330	365	350	308	311	365	365	315	318	365	365	324	323	365	365	334	336	365	365	306	313	365	365
Adult Maint.	365	325	330	365	350	308	311	365	365	315	318	365	365	324	323	365	365	334	336	365	365	306	313	365	365

Number of Annual Degree Days (deg C)

4,151

5,670

5,80

4,926

4,643

4,712

RCR j:\TROA\_03\Fish\Fishsum\FishTempSum4.xls

rev 15-Feb-2004

Date of Truckee River Operating Model Analysis:

10-Feb-04

RCR: 1

\*\*Provisional Data\*\*

17-Feb-2004

Table DSSAMt 19. No. of days recommended temperatures are met for Truckee R. Fishes

Truckee R. 1986wy (extremely wet) - LWSA

Rainbow Trout

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg			
		MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN
Adult Migration	46	39	41	41	33	32	38	46	38	31	36	45	40	31	35	43	40	28	30	38	41	25	28	36	39
Spawning	32	25	27	24	13	18	24	29	19	17	22	29	20	17	21	27	23	14	16	24	28	11	14	22	27
Incubation	62	32	36	40	39	21	27	44	37	19	22	39	37	20	21	36	40	14	16	24	38	11	14	22	37
Rearing	183	178	183	176	173	181	183	183	183	174	177	183	183	175	180	183	183	162	167	183	182	146	158	183	178
Juvenile Maint.	365	356	361	342	335	363	365	365	365	356	359	365	365	357	362	365	365	344	349	365	359	328	340	365	350
Adult Maint.	365	356	361	342	335	363	365	365	365	356	359	365	365	357	362	365	365	344	349	365	359	328	340	365	350

Cui-ui

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg			
		MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN
Adult Migration	150	78	83	46	35	86	88	56	41	73	78	62	45	72	77	62	45	64	64	58	49	61	62	52	46
Spawning	91	64	69	46	35	70	72	54	41	59	64	60	45	58	63	60	45	53	53	56	49	50	51	48	46
Incubation	91	71	71	46	35	78	78	54	41	78	78	60	45	78	78	61	45	75	77	64	49	76	77	65	48
Larvae	62	55	56	47	36	57	58	52	42	57	58	54	46	57	58	54	46	55	58	55	49	54	55	55	48

Lahontan Cutthroat Trout

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg			
		MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN
Adult Migration	136	100	102	114	102	98	102	109	113	96	102	98	107	96	101	96	103	94	100	94	96	92	100	94	94
Spawning	91	59	60	59	44	56	57	61	48	53	57	51	51	54	57	51	50	55	57	49	44	51	57	52	43
Incubation	91	59	60	59	44	56	57	61	48	53	57	51	51	54	57	51	50	55	57	49	44	51	57	52	43
Juvenile Maint.	107	89	95	96	92	92	98	104	96	78	90	107	97	81	91	107	97	75	83	107	98	57	60	107	96
Adult Maint.	365	339	339	332	315	365	365	364	343	358	362	346	336	354	356	343	336	336	339	338	334	330	340	337	333

Mountain Whitefish

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg			
		MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN
Adult Migration	62	38	39	48	51	30	36	41	48	33	36	41	47	40	36	41	46	29	32	41	46	17	14	41	47
Spawning	47	24	22	36	33	25	22	33	47	26	23	33	39	29	27	30	37	29	24	30	36	25	22	29	36
Incubation	78	37	35	47	42	56	53	64	78	57	54	59	55	58	56	53	51	56	51	49	49	55	52	48	48
Incubation	74	74	74	74	72	68	74	74	74	71	74	74	74	72	74	74	74	74	74	74	74	74	74	74	74
Rearing	154	135	136	154	154	135	136	154	154	130	129	151	154	130	130	151	154	126	125	149	154	122	123	148	154
Juvenile Maint.	365	310	308	342	335	315	314	355	365	306	300	340	365	307	301	339	365	301	295	336	359	295	293	334	350
Adult Maint.	365	310	308	342	335	315	314	355	365	306	300	340	365	307	301	339	365	301	295	336	359	295	293	334	350

Brown Trout

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg			
		MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN
Adult Migration	62	38	39	48	51	30	36	41	48	33	36	41	47	40	36	41	46	29	32	41	46	17	14	41	47
Spawning	32	15	13	30	29	10	7	18	32	11	8	18	31	14	12	18	30	14	9	22	31	10	7	21	32
Incubation	78	37	35	47	42	56	53	64	78	57	54	59	55	58	56	53	51	56	51	49	49	55	52	48	48
Incubation	74	74	74	74	72	68	74	74	74	71	74	74	74	72	74	74	74	74	74	74	74	74	74	74	74
Rearing	154	135	136	154	154	135	136	154	154	130	129	151	154	130	130	151	154	126	125	149	154	122	123	148	154
Juvenile Maint.	365	310	308	342	335	315	314	355	365	306	300	340	365	307	301	339	365	301	295	336	359	295	293	334	350
Adult Maint.	365	310	308	342	335	315	314	355	365	306	300	340	365	307	301	339	365	301	295	336	359	295	293	334	350

Number of Annual Degree Days (deg C) 3,383 3,806 3,865 3,874 3,913 3,948

RCR j:\TROA\_03\Fish\FishSum\FishTempSum4 web3

rev 18-Feb-2004

Date of Truckee River Operating Model Analysis: 10-Feb-04

Table DSSAMt 20. No. of days recommended temperatures are met for Truckee R. Fishes

## Truckee R. 1966wy (median) - LWSA

### Rainbow Trout

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg			
		Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN
Adult Migration	46	36	41	46	46	30	35	46	46	21	28	45	46	21	28	40	46	7	1	31	43	2	0	27	42
Spawning	32	22	27	32	16	18	21	32	28	12	14	31	32	12	14	26	32	4	1	17	29	1	0	13	28
Incubation	62	24	27	55	46	19	21	44	54	12	14	31	47	12	14	26	37	4	1	17	29	1	0	13	28
Rearing	183	182	183	183	183	183	183	183	183	181	183	183	183	180	183	183	183	161	162	183	183	139	141	183	183
Juvenile Maint.	365	364	365	360	341	365	365	365	365	363	365	365	365	362	365	365	365	343	344	365	365	321	323	365	365
Adult Maint.	365	364	365	360	341	365	365	365	365	363	365	365	365	362	365	365	365	343	344	365	365	321	323	365	365

### Cui-ui

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg			
		Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN
Adult Migration	150	87	90	87	57	89	95	95	86	77	82	84	90	76	79	82	89	59	70	71	79	42	41	68	82
Spawning	91	73	76	85	57	70	75	87	85	62	66	77	88	61	64	76	87	46	55	67	77	28	23	63	80
Incubation	91	86	88	88	57	85	88	91	85	82	87	90	90	62	86	90	91	71	75	87	91	62	65	82	91
Larvae	62	57	58	62	55	56	58	62	62	53	57	60	62	52	56	60	62	41	45	57	62	32	35	52	62

### Lahontan Cutthroat Trout

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg			
		Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN
Adult Migration	136	57	63	94	96	57	59	97	103	44	50	73	90	42	48	64	84	37	32	41	70	35	34	40	67
Spawning	91	40	44	76	75	29	30	70	82	23	23	58	75	22	23	54	69	10	12	29	61	6	0	23	60
Incubation	91	40	44	76	75	29	30	70	82	23	23	58	75	22	23	54	69	10	12	29	61	6	0	23	60
Juvenile Maint.	107	102	107	107	101	103	107	107	107	94	99	107	107	94	97	107	107	70	71	107	107	47	50	107	107
Adult Maint.	365	356	356	327	302	365	365	364	352	365	365	365	342	364	364	352	339	346	349	347	332	325	329	346	324

### Mountain Whitefish

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg			
		Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN
Adult Migration	62	22	17	42	62	20	17	19	45	18	16	18	31	18	16	17	30	15	15	16	19	9	7	16	22
Spawning	47	29	31	46	47	24	17	45	46	31	27	39	45	31	28	38	45	30	24	34	44	28	22	33	44
Incubation	78	60	62	75	68	55	48	76	77	62	58	70	73	62	59	69	73	61	55	62	67	59	53	59	65
Incubation	74	64	68	53	28	65	70	74	74	71	73	74	67	71	74	74	66	71	74	71	63	71	73	71	58
Rearing	154	137	141	154	153	135	139	154	154	132	135	149	154	132	135	145	154	110	107	140	154	102	104	137	154
Juvenile Maint.	365	331	332	360	341	330	330	365	365	323	323	346	365	322	323	338	365	290	288	328	365	271	276	325	365
Adult Maint.	365	331	332	360	341	330	330	365	365	323	323	346	365	322	323	338	365	290	288	328	365	271	276	325	365

### Brown Trout

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg				7-day moving avg			
		Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN	Inst. MAX	MAX	MEAN	MIN
Adult Migration	62	22	17	42	62	20	17	19	45	18	16	18	31	18	16	17	30	15	15	16	19	9	7	16	22
Spawning	32	14	16	31	32	9	5	30	31	16	12	24	30	16	13	23	30	15	9	19	29	13	7	18	29
Incubation	78	60	62	75	68	55	48	76	77	62	58	70	73	62	59	69	73	61	55	62	67	59	53	59	65
Incubation	74	64	68	53	28	65	70	74	74	71	73	74	67	71	74	74	66	71	74	71	63	71	73	71	58
Rearing	154	153	154	154	153	154	154	154	154	153	154	154	154	153	154	154	154	147	147	154	154	131	135	154	154
Juvenile Maint.	365	364	365	360	341	365	365	365	365	363	365	365	365	362	365	365	365	343	344	365	365	321	323	365	365
Adult Maint.	365	364	365	360	341	365	365	365	365	363	365	365	365	362	365	365	365	343	344	365	365	321	323	365	365

31

Number of Annual Degree Days (deg C)

3,469

3,969

4,140

4,177

4,347

4,416

RCR j:\TRQA\_03\Fish\FishSum\FishTempSum4web3

rev 18-Feb-2004

Date of Truckee River Operating Model Analysis:

10-Feb-04

RCR, 1

\*\*Provisional Data\*\*

17-Feb-2004

Table DSSAMt 21. No. of days recommended temperatures are met for Truckee R. Fishes

# Truckee R. 1992wy (extremely dry) - LWSA

## Rainbow Trout

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg			
		MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN
Adult Migration	46	3	0	12	33	1	0	1	19	0	0	2	11	0	0	2	11	1	0	3	12	0	0	3	25
Spawning	32	2	0	6	19	0	0	0	11	0	0	0	4	0	0	0	4	1	0	0	5	0	0	0	13
Incubation	62	2	0	11	42	0	0	0	11	0	0	0	9	0	0	0	9	1	0	0	11	0	0	0	28
Rearing	183	143	149	183	183	127	129	183	183	133	137	183	183	142	141	183	183	153	154	183	183	129	132	183	183
Juvenile Maint	365	325	331	365	352	309	311	365	365	315	319	365	365	324	323	365	365	334	336	365	365	306	314	365	365
Adult Maint	365	325	331	365	352	309	311	365	365	315	319	365	365	324	323	365	365	334	336	365	365	306	314	365	365

## Cui-ui

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg			
		MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN
Adult Migration	150	61	60	66	68	83	85	83	95	65	66	71	78	63	65	67	66	63	65	59	67	53	60	68	73
Spawning	91	29	28	63	68	26	28	39	62	29	30	37	61	28	30	37	58	29	31	40	61	19	24	41	68
Incubation	91	57	61	87	74	49	50	69	81	48	51	71	91	50	52	69	88	50	51	69	91	38	36	71	91
Larvae	62	28	31	58	62	19	20	39	52	18	21	42	62	21	22	40	59	21	21	40	62	9	6	42	62

## Lahontan Cutthroat Trout

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg			
		MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN
Adult Migration	136	52	59	66	85	43	42	70	73	47	53	71	74	46	51	72	67	50	53	65	67	37	35	68	72
Spawning	91	14	24	31	65	9	11	26	32	15	15	28	31	14	14	28	32	14	13	29	34	8	7	28	41
Incubation	91	14	24	31	65	9	11	26	32	15	15	28	31	14	14	28	32	14	13	29	34	8	7	28	41
Juvenile Maint	107	65	66	107	107	32	34	92	107	56	59	96	107	59	61	95	107	66	68	107	107	53	54	107	107
Adult Maint.	365	331	335	329	309	314	319	365	365	323	324	365	365	333	331	365	365	342	345	345	326	311	318	364	338

## Mountain Whitefish

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg			
		MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN
Adult Migration	62	11	8	11	42	1	0	3	5	9	6	7	9	9	6	7	9	10	7	8	10	9	6	8	11
Spawning	47	30	27	35	39	2	0	1	13	25	19	25	29	26	22	27	30	30	28	35	35	25	21	31	36
Incubation	78	61	58	59	44	12	10	30	44	56	50	56	60	57	53	58	61	61	59	66	60	56	52	62	64
Incubation	74	59	63	63	52	25	24	47	65	55	55	65	74	56	56	69	74	60	63	66	57	54	55	71	66
Rearing	154	102	102	150	154	90	92	115	152	94	95	123	152	96	98	123	149	94	98	127	152	80	78	132	154
Juvenile Maint	365	288	286	353	352	226	224	291	349	244	239	311	354	256	254	311	347	273	280	316	354	231	221	323	365
Adult Maint.	365	288	286	353	352	226	224	291	349	244	239	311	354	256	254	311	347	273	280	316	354	231	221	323	365

## Brown Trout

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg			
		MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN
Adult Migration	62	11	8	11	42	1	0	3	5	9	6	7	9	9	6	7	9	10	7	8	10	9	6	8	11
Spawning	32	15	12	20	24	0	0	0	0	10	6	10	14	11	8	12	15	15	13	20	20	10	7	16	21
Incubation	78	61	58	59	44	12	10	30	44	56	50	56	60	57	53	58	61	61	59	66	60	56	52	62	64
Incubation	74	59	63	63	52	25	24	47	65	55	55	65	74	56	56	69	74	60	63	66	57	54	55	71	66
Rearing	154	128	141	154	154	120	123	154	154	122	130	154	154	129	134	154	154	137	143	154	154	114	125	154	154
Juvenile Maint	365	325	331	365	352	309	311	365	365	315	319	365	365	324	323	365	365	334	336	365	365	306	314	365	365
Adult Maint.	365	325	331	365	352	309	311	365	365	315	319	365	365	324	323	365	365	334	336	365	365	306	314	365	365

Number of Annual Degree Days (deg C)

4,155

5,675

5,076

4,930

4,643

4,712

RCR j:\TROA\_03\Fish\FishSum\FishTempSum4.tbl3

rev 18-Feb-2004

Date of Truckee River Operating Model Analysis:

10-Feb-04

RCR, 1

\*\*Provisional Data\*\*

17-Feb-2004

Table DSSAmT 22. No. of days recommended temperatures are met for Truckee R. Fishes

## Truckee R. 1986wy (extremely wet) - TROA

### Rainbow Trout

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg			
		MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN
Adult Migration	46	36	41	46	46	30	35	46	46	21	28	45	46	21	28	40	48	7	1	31	43	2	0	27	42
Spawning	32	22	27	32	16	18	21	32	28	12	14	31	32	12	14	26	32	4	1	17	29	1	0	13	28
Incubation	62	24	27	55	46	19	21	44	54	12	14	31	47	12	14	26	37	4	1	17	29	1	0	13	28
Rearing	183	182	183	183	183	183	183	183	183	181	183	183	183	180	183	183	183	161	162	183	183	139	141	183	183
Juvenile Maint.	365	364	365	360	341	365	365	365	365	363	365	365	365	362	365	365	365	343	344	365	365	321	323	365	365
Adult Maint.	365	364	365	360	341	365	365	365	365	363	365	365	365	362	365	365	365	343	344	365	365	321	323	365	365

### Cui-ui

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg			
		MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN
Adult Migration	150	87	90	87	57	89	95	95	86	77	82	84	90	76	79	82	89	59	70	71	79	43	41	68	82
Spawning	91	73	76	85	57	70	75	87	85	62	66	77	88	61	64	76	87	46	55	67	77	29	23	63	80
Incubation	91	86	88	88	57	85	88	91	85	82	87	90	90	82	86	90	91	71	75	87	91	62	65	82	91
Larvae	62	57	58	62	55	56	56	62	62	53	57	60	62	52	56	60	62	41	45	57	62	32	35	52	62

### Lahontan Cutthroat Trout

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg			
		MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN
Adult Migration	136	58	63	94	96	59	80	97	102	44	50	73	90	42	48	65	84	37	35	42	70	35	34	41	67
Spawning	91	40	46	76	74	32	31	70	82	24	23	58	75	22	23	54	69	11	12	29	61	6	0	23	60
Incubation	91	40	46	76	74	32	31	70	82	24	23	58	75	22	23	54	69	11	12	29	61	6	0	23	60
Juvenile Maint.	107	102	107	107	101	103	107	107	107	94	100	107	107	94	97	107	107	71	71	107	107	47	50	107	107
Adult Maint.	365	356	356	327	301	365	365	364	352	365	365	355	342	364	364	352	339	346	349	347	332	324	328	344	324

### Mountain Whitefish

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg			
		MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN
Adult Migration	62	22	17	42	62	20	17	19	45	18	16	18	31	18	16	17	30	15	15	16	20	9	7	16	22
Spawning	47	29	31	46	47	24	18	45	46	31	27	39	45	31	28	38	45	31	24	34	44	28	22	33	44
Incubation	78	60	62	75	88	55	49	76	77	62	58	70	73	62	59	69	73	62	55	62	67	59	53	59	65
Incubation	74	64	68	53	28	65	70	74	74	71	73	74	67	71	74	73	66	71	74	71	63	71	73	70	58
Rearing	154	137	141	154	153	135	139	154	154	132	135	149	154	132	135	145	154	110	107	140	154	102	104	137	154
Juvenile Maint.	365	331	332	360	341	330	330	365	365	323	323	346	365	322	323	338	365	290	288	328	365	271	276	325	365
Adult Maint.	365	331	332	360	341	330	330	365	365	323	323	346	365	322	323	338	365	290	288	328	365	271	276	325	365

### Brown Trout

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg			
		MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN
Adult Migration	62	22	17	42	62	20	17	19	45	18	16	18	31	18	16	17	30	15	15	16	20	9	7	16	22
Spawning	32	14	16	31	32	9	5	30	31	16	12	24	30	16	13	23	30	16	9	19	29	13	7	18	29
Incubation	78	60	62	75	88	55	49	76	77	62	58	70	73	62	59	69	73	62	55	62	67	59	53	59	65
Incubation	74	64	68	53	28	65	70	74	74	71	73	74	67	71	74	73	66	71	74	71	63	71	73	70	58
Rearing	154	153	154	154	153	154	154	154	154	153	154	154	154	153	154	154	154	147	147	154	154	131	135	154	154
Juvenile Maint.	365	364	365	360	341	365	365	365	365	363	365	365	365	362	365	365	365	343	344	365	365	321	323	365	365
Adult Maint.	365	364	365	360	341	365	365	365	365	363	365	365	365	362	365	365	365	343	344	365	365	321	323	365	365

3

Number of Annual Degree Days (deg C)

3,466

3,963

4,136

4,173

4,344

4,413

RCR j:\TROA\_03\Fish\Fishsum\FishTempSum4.xls

rev 18-Feb-2004

Date of Truckee River Operating Model Analysis:

10-Feb-04

RCR 1

Provisional Data

17-Feb-2004

Table DSSAMt 23. No. of days recommended temperatures are met for Truckee R. Fishes

## Truckee R. 1966wy (median) - TROA

### Rainbow Trout

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg			
		MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN
Adult Migration	46	25	28	44	46	16	20	42	46	9	0	31	44	8	0	29	44	0	0	13	34	0	0	3	34
Spawning	32	13	14	30	21	8	10	28	32	5	0	17	30	5	0	15	30	0	0	7	20	0	0	1	20
Incubation	62	13	14	32	48	8	10	28	45	5	0	17	30	5	0	15	30	0	0	7	20	0	0	1	20
Rearing	183	182	183	183	182	183	183	183	183	179	183	183	183	177	183	183	183	157	161	163	183	125	128	183	183
Juvenile Maint.	365	364	365	360	341	365	365	365	365	361	365	365	365	359	365	365	365	339	343	365	365	307	310	365	365
Adult Maint.	365	364	365	360	341	365	365	365	365	361	365	365	365	359	365	365	365	339	343	365	365	307	310	365	365

### Cui-ui

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg			
		MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN
Adult Migration	150	79	85	78	63	82	86	87	81	66	74	81	80	65	72	77	78	40	42	64	76	36	38	61	73
Spawning	91	64	70	76	63	61	64	74	79	49	56	72	77	49	55	68	75	27	26	59	73	22	21	56	70
Incubation	91	77	79	98	63	76	78	91	85	73	74	84	91	73	74	82	91	62	64	80	89	53	58	78	91
Larvae	62	48	49	62	59	47	48	62	62	43	44	54	62	43	44	52	62	32	34	50	60	23	28	48	62

### Lahontan Cutthroat Trout

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg			
		MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN
Adult Migration	136	48	49	84	85	51	54	82	98	43	49	55	80	41	47	52	70	34	34	40	53	36	36	39	54
Spawning	91	26	24	70	67	19	19	58	73	16	19	35	60	16	19	28	58	10	11	22	45	5	0	22	45
Incubation	91	26	24	70	67	19	19	58	73	16	19	35	60	16	19	28	58	10	11	22	45	5	0	22	45
Juvenile Maint.	107	100	107	107	101	100	107	107	107	85	90	107	107	81	88	107	107	67	66	107	107	43	39	107	107
Adult Maint.	365	356	356	325	292	365	365	365	357	362	365	360	346	362	365	356	345	345	351	350	331	319	320	347	320

### Mountain Whitefish

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg			
		MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN
Adult Migration	62	28	23	39	62	25	19	28	45	23	18	26	34	23	18	26	32	20	17	20	26	16	16	18	26
Spawning	47	33	31	45	42	26	26	46	47	33	29	45	47	33	31	45	46	32	27	44	45	29	22	39	45
Incubation	78	64	62	75	61	57	57	77	78	64	60	76	78	64	62	76	77	63	58	73	69	60	53	66	65
Incubation	74	63	66	54	30	64	69	71	74	70	72	74	68	71	72	74	67	71	74	72	64	71	73	71	59
Rearing	154	130	133	154	154	128	131	154	154	114	117	146	154	113	116	141	154	104	102	132	154	94	93	130	154
Juvenile Maint.	365	324	324	360	341	322	321	365	365	303	304	338	365	302	303	332	365	283	282	320	362	263	264	317	365
Adult Maint.	365	324	324	360	341	322	321	365	365	303	304	338	365	302	303	332	365	283	282	320	362	263	264	317	365

### Brown Trout

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg			
		MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN
Adult Migration	62	28	23	39	62	25	19	28	45	23	18	26	34	23	18	26	32	20	17	20	26	16	16	18	26
Spawning	32	18	16	32	32	11	11	31	32	18	14	30	32	18	16	30	31	17	12	29	30	14	7	24	30
Incubation	78	64	62	75	61	57	57	77	78	64	60	76	78	64	62	76	77	63	58	73	69	60	53	66	65
Incubation	74	63	66	54	30	64	69	71	74	70	72	74	68	71	72	74	67	71	74	72	64	71	73	71	59
Rearing	154	153	154	154	154	154	154	154	154	153	154	154	154	152	154	154	154	143	146	154	154	118	123	154	154
Juvenile Maint.	365	364	365	360	341	365	365	365	365	361	365	365	365	359	365	365	365	339	343	365	365	307	310	365	365
Adult Maint.	365	364	365	360	341	365	365	365	365	361	365	365	365	359	365	365	365	339	343	365	365	307	310	365	365

Number of Annual Degree Days (deg C)

3,549

4,088

4,246

4,281

4,405

4,455

RGR j:\TROA\_03\Fish\Fishsum\FishTempSum4.wb3

rev 16-Feb-2004

Date of Truckee River Operating Model Analysis:

10-Feb-04

RGR: 1

"Provisional Data"

17-Feb-2004



Table DSSAMT 24. No. of days recommended temperatures are met for Truckee R. Fishes

Truckee R. 1992wy (extremely dry) - TROA

Rainbow Trout

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg			
		MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN
Adult Migration	46	3	0	12	33	1	0	1	19	0	0	2	11	0	0	2	11	1	0	3	13	0	0	3	24
Spawning	32	2	0	6	19	0	0	0	11	0	0	0	4	0	0	0	4	1	0	0	6	0	0	0	12
Incubation	62	2	0	11	40	0	0	0	14	0	0	0	9	0	0	0	8	1	0	0	12	0	0	0	24
Rearing	183	156	163	183	183	147	154	183	163	141	141	183	183	147	153	183	183	163	156	183	183	130	133	183	183
Juvenile Maint.	365	338	345	365	351	329	336	365	365	323	323	365	365	329	335	365	365	334	338	365	365	307	315	365	365
Adult Maint.	365	338	345	365	351	329	336	365	365	323	323	365	365	329	335	365	365	334	338	365	365	307	315	365	365

Cui-ui

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg			
		MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN
Adult Migration	150	61	60	66	66	85	87	87	98	65	66	71	78	63	65	68	69	63	65	59	67	53	60	69	74
Spawning	91	29	28	63	66	26	26	40	64	29	30	37	61	28	30	37	58	29	31	40	61	19	24	42	68
Incubation	91	59	62	86	74	51	54	71	87	49	51	71	91	51	52	69	87	50	51	69	91	39	36	71	91
Larvae	62	30	32	57	62	22	24	42	58	20	21	42	62	22	22	40	58	21	21	40	62	9	6	42	62

Lahontan Cutthroat Trout

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg			
		MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN
Adult Migration	136	52	59	68	85	43	42	70	73	47	53	71	74	46	51	72	68	50	54	65	68	37	35	67	71
Spawning	91	16	24	32	65	10	10	27	32	15	16	28	31	14	14	28	32	14	13	29	34	8	7	28	40
Incubation	91	16	24	32	65	10	10	27	32	15	16	28	31	14	14	28	32	14	13	29	34	8	7	28	40
Juvenile Maint.	107	72	76	107	107	68	68	107	107	60	61	98	107	64	65	97	107	66	68	106	107	53	55	105	107
Adult Maint.	365	340	346	325	309	341	349	365	365	329	336	365	365	338	343	365	365	345	347	345	324	314	318	364	339

Mountain Whitefish

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg			
		MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN
Adult Migration	62	10	8	10	16	7	4	6	7	9	6	7	9	10	6	7	9	10	7	8	9	9	6	8	10
Spawning	47	30	26	36	38	2	0	1	8	26	20	26	29	28	23	27	31	30	27	34	35	27	22	31	36
Incubation	78	61	57	60	43	13	9	32	39	57	51	57	60	59	54	58	62	61	58	65	60	58	53	62	64
Incubation	74	59	63	63	52	23	24	47	64	55	55	65	74	56	56	68	74	60	63	66	57	54	55	71	66
Rearing	154	104	105	151	154	98	99	128	154	95	97	124	152	97	97	124	149	96	97	128	152	81	79	130	154
Juvenile Maint.	365	289	290	349	351	253	245	317	360	251	248	312	351	263	264	312	347	277	276	317	354	232	224	319	363
Adult Maint.	365	289	290	349	351	253	245	317	360	251	248	312	351	263	264	312	347	277	276	317	354	232	224	319	363

Brown Trout

Life Stage	Total # Days	East McCarran 96.7 km				Lockwood 106.5 km				Clark 124.9 km				Painted Rock 131.6 km				Dead Ox 171.7 km				Marble Bluff 187.0 km			
		Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg				Inst. 7-day moving avg			
		MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN	MAX	MAX	MEAN	MIN
Adult Migration	62	10	8	10	16	7	4	6	7	9	6	7	9	10	6	7	9	10	7	8	9	9	6	8	10
Spawning	32	15	11	21	23	0	0	1	8	11	7	11	14	13	9	12	16	15	12	19	20	12	8	16	21
Incubation	78	61	57	60	43	13	9	32	39	57	51	57	60	59	54	58	62	61	58	65	60	58	53	62	64
Incubation	74	59	63	63	52	23	24	47	64	55	55	65	74	56	56	68	74	60	63	66	57	54	55	71	66
Rearing	154	138	150	154	154	132	143	154	154	128	134	154	154	134	143	154	154	137	144	154	154	115	126	154	154
Juvenile Maint.	365	338	345	365	351	329	336	365	365	323	323	365	365	329	335	365	365	334	338	365	365	307	315	365	365
Adult Maint.	365	338	345	365	351	329	336	365	365	323	323	365	365	329	335	365	365	334	338	365	365	307	315	365	365

Number of Annual Degree Days (deg C) 4,208 5,485 5,060 4,925 4,653 4,748

ROR: 1 TROA\_03/Fish/Fishsum/FishTempSum4.wb3

rev 18-Feb-2004

Date of Truckee River Operating Model Analysis: 10-Feb-04

## REFERENCES

- Bender, Merlynn D., February 1995, "Summary of Information for Estimating Water Quality Conditions Downstream of Lake Tahoe for the TROA RDEIS/RDEIR," Technical Service Center, U.S. Bureau of Reclamation, Special Report, Denver, Colorado.
- Bureau of Reclamation, May 1986, "Draft Environmental Impact Statement for the Newlands Project Proposed Operating Criteria and Procedures." Prepared by URS Corporation for the Mid-Pacific Regional Office, Sacramento, California.
- Bureau of Reclamation, December 1987, "Final Environmental Impact Statement for the Newlands Project Proposed Operating Criteria and Procedures." Prepared by URS Corporation for the Mid-Pacific Regional Office, Sacramento, California.
- Chen, Carl W., and Laura H.Z. Weintraub, updated March 2002, "Adaptation of WARMF to the Truckee River Basin of California and Nevada." Prepared by Systech Engineering Inc. for City of Reno, NV, City of Sparks, NV, and Washoe County, NV.
- Hallock, Robert J., and Linda L. Hallock, Editors, U.S. Fish and Wildlife Service, 1993, "Detailed Study of Irrigation Drainage in and near Wildlife Management Areas, West-Central Nevada, 1987-90, Part B. Effect on Biota in Stillwater and Fernley Wildlife Management Areas and Other Nearby Wetlands," U.S. Geological Survey, Water- Resources Investigations Report 92-4042B, Carson City, Nevada.
- Hoffman, Ray J., Robert J. Hallock, Timothy G. Rowe, Michael S. Lico, Howard L. Burge, and Steven P. Thompson, 1990, "Reconnaissance Investigation of Water Quality, Bottom Sediment, and Biota Associated With Irrigation Drainage in and near Stillwater Wildlife Management Area, Churchill County, Nevada, 1986-87," U.S. Geological Survey, Water-Resources Investigations Report 89-4105, Carson City, Nevada.
- Lahontan Regional Water Quality Control Board, 1995, "Water Quality Plan for the Lahontan Region," Sacramento, California.
- Lebo, Martin E., John E. Reuter, Cathryn L. Rhodes, and Charles R. Goldman, "Pyramid Lake, Nevada Water Quality Study 1989-1993, Volume 1. Limnological Data," Division of Environmental Studies, University of California, Davis, California, October 1993.
- Lebo, Martin E., John E. Reuter, Cathryn L. Rhodes, and Charles R. Goldman, "Pyramid Lake, Nevada Water Quality Study 1989-1993, Volume 2. Limnological Description," Division of Environmental Studies, University of California, Davis, California, October 1993.
- Lebo, Martin E., John E. Reuter, and Charles R. Goldman, "Pyramid Lake, Nevada Water Quality Study 1989-1993, Volume 3. Nutrient Budgets," Division of Environmental Studies, University of California, Davis, California, October 1993.

Lebo, Martin E., John E. Reuter, Cathryn L. Rhodes, and Charles R. Goldman, "Pyramid Lake, Nevada Water Quality Study 1989-1993, Volume 4. Modeling Studies," Division of Environmental Studies, University of California, Davis, California, October 1994.

Lico, Michael S., 1992, "Detailed Study of Irrigation Drainage in and near Wildlife Management Areas, West-Central Nevada, 1987-90, Part A. Water Quality, Sediment Composition, and Hydrogeochemical Processes in Stillwater and Fernley Wildlife Management Areas," U.S. Geological Survey, Water Resources Investigations Report 92-4024A, Carson City, Nevada.

Marshack, Jon B., 1993, "A Compilation of Water Quality Goals," California Regional Water Quality Control Board, Central Valley Region, Sacramento, California.

Marshack, Jon B., August 2003, "A Compilation of Water Quality Goals," Regional Water Quality Control Board, Central Valley Region, California Environmental Protection Agency.

Rowe, Timothy G., Michael S. Lico, Robert J. Hallock, Ann S. Maest, Ann S., Ray J. Hoffman, 1991, "Physical, Chemical, and Biological Data for Detailed Study of Irrigation Drainage in and near Stillwater, Fernley, and Humboldt Wildlife Management Areas and Carson Lake, West-Central Nevada, 1987-89," U.S. Geological Survey, Open-File Report 91-185, Carson City, Nevada.

Seiler, Ralph L., Geoffrey A. Ekechukwu, and Robert J. Hallock, 1993, "Reconnaissance Investigation of Water Quality, Bottom Sediment, and Biota Associated with Irrigation Drainage in and near Humboldt Wildlife Management Area, Churchill and Pershing Counties, Nevada, 1990-91," U.S. Geological Survey, Water Resources Investigations Report 93-4072, Carson City, Nevada.

New Reference: - need to check where listed in document

U.S. Department of the Interior, Bureau of Indian Affairs, October 2002, "Truckee River Water Quality Settlement Agreement - Federal Water Rights Acquisition Program, Final Environmental Impact Statement," California and Nevada, Western Regional Office of the Bureau of Indian Affairs, Phoenix, Arizona.

New Reference: - need to check where listed in document

U.S. Department of the Interior, Bureau of Indian Affairs, October 2002, "Truckee River Water Quality Settlement Agreement - Federal Water Rights Acquisition Program, Final Environmental Impact Statement," California and Nevada, Western Regional Office of the Bureau of Indian Affairs, Phoenix, Arizona.

Rapid Creek Research, Inc. Modeling References (In Preparation).

- Caupp, C.L., J.T. Brock, and H.M. Runke, 2004. "Application of the Dynamic Stream Simulation and Assessment Model (DSSAMt) to the Truckee River, Nevada for Truckee River Operating Agreement (TROA) RDEIS/RDEIR: Model Formulation and Overview," Technical Report No. RCR04-1.0. Submitted to U.S. Bureau of Reclamation. Carson City, Nevada. Rapid Creek Research, Inc., Boise, Idaho. (in preparation).
- Brock, J.T. and C.L. Caupp. 2004 . "Calibration of DSSAMt Water Quality Model - Truckee River, Nevada for Truckee River Operating Agreement (TROA) RDEIS/RDEIR: Model Performance and Sensitivity of Simulation Results for 1986, 1989, 1991, 1992, and 1993." Technical Report No. RCR04-2.0. Submitted to U.S. Bureau of Reclamation. Carson City, Nevada. Rapid Creek Research, Inc., Boise, Idaho. (in preparation).
- Brock, J.T. and C.L. Caupp. 2004 . "Application of DSSAMt Water Quality Model - Truckee River, Nevada for Truckee River Operating Agreement (TROA) RDEIS/RDEIR: Simulated River Temperatures for Current Condition." Technical Report No. RCR04-3.0. Submitted to U.S. Bureau of Reclamation. Carson City, Nevada. Rapid Creek Research, Inc., Boise, Idaho. (in preparation).
- Brock, J.T. and C.L. Caupp. 2004 . "Application of DSSAMt Water Quality Model - Truckee River, Nevada for Truckee River Operating Agreement (TROA) RDEIS/RDEIR: Simulated Water Quality for Current Condition." Technical Report No. RCR04-4.0. Submitted to U.S. Bureau of Reclamation. Carson City, Nevada. Rapid Creek Research, Inc., Boise, Idaho. (in preparation).
- Brock, J.T. and C.L. Caupp. 2004 . "Application of DSSAMt Water Quality Model - Truckee River, Nevada for Truckee River Operating Agreement (TROA) RDEIS/RDEIR: Simulated River Temperatures for No Action." Technical Report No. RCR04-5.0. Submitted to U.S. Bureau of Reclamation. Carson City, Nevada. Rapid Creek Research, Inc., Boise, Idaho. (in preparation).
- Brock, J.T. and C.L. Caupp. 2004 . "Application of DSSAMt Water Quality Model - Truckee River, Nevada for Truckee River Operating Agreement (TROA) RDEIS/RDEIR: Simulated Water Quality for No Action." Technical Report No. RCR04-6.0. Submitted to U.S. Bureau of Reclamation. Carson City, Nevada. Rapid Creek Research, Inc., Boise, Idaho. (in preparation).

- Brock, J.T. and C.L. Caupp. 2004 . “Application of DSSAMt Water Quality Model - Truckee River, Nevada for Truckee River Operating Agreement (TROA) RDEIS/RDEIR: Simulated River Temperatures for TROA.” Technical Report No. RCR04-7.0. Submitted to U.S. Bureau of Reclamation. Carson City, Nevada. Rapid Creek Research, Inc., Boise, Idaho.  
(in preparation).
- Brock, J.T. and C.L. Caupp. 2004 . “Application of DSSAMt Water Quality Model - Truckee River, Nevada for Truckee River Operating Agreement (TROA) RDEIS/RDEIR: Simulated Water Quality for TROA.” Technical Report No. RCR04-8.0. Submitted to U.S. Bureau of Reclamation. Carson City, Nevada. Rapid Creek Research, Inc., Boise, Idaho.  
(in preparation).
- Brock, J.T. and C.L. Caupp. 2004 . “Application of DSSAMt Water Quality Model - Truckee River, Nevada for Truckee River Operating Agreement (TROA) RDEIS/RDEIR: Simulated River Temperatures for LWSA.” Technical Report No. RCR04-9.0. Submitted to U.S. Bureau of Reclamation. Carson City, Nevada. Rapid Creek Research, Inc., Boise, Idaho.  
(in preparation).
- Brock, J.T. and C.L. Caupp. 2004 . “Application of DSSAMt Water Quality Model - Truckee River, Nevada for Truckee River Operating Agreement (TROA) RDEIS/RDEIR: Simulated Water Quality for LWSA” Technical Report No. RCR04-10.0. Submitted to U.S. Bureau of Reclamation. Carson City, Nevada. Rapid Creek Research, Inc., Boise, Idaho.  
(in preparation).

# **Attachment to Water Quality Appendix**

## **Beneficial Uses of Surface Waters**

Beneficial uses of surface waters of the Lahontan Region are listed at:  
<http://www.swrcb.ca.gov/rwqcb6/BasinPlan/Index.htm> and are attached.

A summary of beneficial uses for water bodies identified in the Nevada Administrative Code are listed at: <http://ndep.nv.gov/bwqp/file/uses> and are attached.

**TABLE 2-1. BENEFICIAL USES OF SURFACE WATERS OF THE LAHONTAN REGION**

[illegible]

**TABLE 2-1. BENEFICIAL USES OF SURFACE WATERS OF THE LAHONTAN REGION**

[illegible]



**TABLE 2-1. BENEFICIAL USES OF SURFACE WATERS OF THE LAHONTAN REGION**

[illegible]

**TABLE 2-1. BENEFICIAL USES OF SURFACE WATERS OF THE LAHONTAN REGION**

[illegible]

**TABLE 2-1. BENEFICIAL USES OF SURFACE WATERS OF THE LAHONTAN REGION**

[illegible]

**TABLE 2-1. BENEFICIAL USES OF SURFACE WATERS OF THE LAHONTAN REGION**

[illegible]

**TABLE 2-1. BENEFICIAL USES OF SURFACE WATERS OF THE LAHONTAN REGION**

[illegible]

**TABLE 2-1. BENEFICIAL USES OF SURFACE WATERS OF THE LAHONTAN REGION**

[illegible]

TABLE 2-1. BENEFICIAL USES OF SURFACE WATERS OF THE LAHONTAN REGION

HU No.	HYDROLOGIC UNIT/SUBUNIT DRAINAGE FEATURE	WATERBODY CLASS MODIFIER	BENEFICIAL USES																RECEIVING WATER
			IR	AR	SR	DR	FR	PR	AV	SC-1	SC-2	SC-3	SC-4	SC-5	SC-6	SC-7	SC-8		
607.20	CAMERO HYDROLOGIC AREA																		
	MINOR SURFACE WATERS		X				X			X	X			X		X			
	MINOR WETLANDS	SPRING/DEEP/EMERGENT/MARSHES	X				X	X		X	X			X	X		X	X	
608.00	FACE TRACK HYDROLOGIC UNIT																		
	MINOR SURFACE WATERS		X				X			X	X			X	X	X	X		
	MINOR WETLANDS	SPRING/DEEP/EMERGENT/MARSHES	X				X	X		X	X			X	X		X	X	
608.10	TEASSETT JUNCTION HYDROLOGIC AREA																		
	MINOR SURFACE WATERS		X				X			X	X			X					
	MINOR WETLANDS	SPRING/DEEP/EMERGENT/MARSHES	X				X	X		X	X			X	X		X	X	
608.20	HINDEN VALLEY HYDROLOGIC AREA																		
	MINOR SURFACE WATERS		X				X			X	X			X					
	MINOR WETLANDS	SPRING/DEEP/EMERGENT/MARSHES	X				X	X		X	X			X	X		X	X	
608.30	SEIDA HYDROLOGIC AREA																		
	MINOR SURFACE WATERS		X				X			X	X			X					
	MINOR WETLANDS	SPRING/DEEP/EMERGENT/MARSHES	X				X	X		X	X			X	X		X	X	
608.40	RAID FLAT HYDROLOGIC AREA																		
	MINOR SURFACE WATERS		X				X			X	X			X					
	MINOR WETLANDS	SPRING/DEEP/EMERGENT/MARSHES	X				X	X		X	X			X	X		X	X	
609.00	ANABOGA HYDROLOGIC UNIT																		
	1.0000000000000000	WETLANDS	X							X	X			X	X	X	X		
	2.0000000000000000	WETLANDS	X							X	X			X	X	X	X		
	ANABOGA RIVER WETLANDS	WETLANDS	X	X						X	X			X	X	X	X		
	ANABOGA RIVER	INTERMITTENT STREAM	X	X						X	X			X	X	X	X		
	ANABOGA RIVER	PERMANENT STREAM	X	X						X	X			X	X	X	X		
	ANABOGA RIVER	PERMANENT STREAM	X	X						X	X			X	X	X	X		
	ANABOGA RIVER	PERMANENT STREAM	X	X						X	X			X	X	X	X		
	ANABOGA RIVER	PERMANENT STREAM	X	X						X	X			X	X	X	X		
	ANABOGA RIVER	PERMANENT STREAM	X	X						X	X			X	X	X	X		
	ANABOGA RIVER	PERMANENT STREAM	X	X						X	X			X	X	X	X		
	ANABOGA RIVER	PERMANENT STREAM	X	X						X	X			X	X	X	X		
609.10	DEATH VALLEY HYDROLOGIC AREA																		
	MINOR SURFACE WATERS		X				X			X	X			X		X			
	MINOR WETLANDS	SPRING/DEEP/EMERGENT/MARSHES	X				X	X		X	X			X	X		X	X	
	STOVEPIPE WELLS HYDROLOGIC SUBAREA																		
	UNDEVELOPED	SPRING/DEEP/EMERGENT/MARSHES	X	X			X	X		X	X			X	X	X	X		
	ANABOGA RIVER	SPRING/DEEP/EMERGENT/MARSHES	X	X			X	X		X	X			X	X	X	X		
	ANABOGA RIVER	SPRING/DEEP/EMERGENT/MARSHES	X	X			X	X		X	X			X	X	X	X		
	ANABOGA RIVER	SPRING/DEEP/EMERGENT/MARSHES	X	X			X	X		X	X			X	X	X	X		
	ANABOGA RIVER	SPRING/DEEP/EMERGENT/MARSHES	X	X			X	X		X	X			X	X	X	X		
	ANABOGA RIVER	SPRING/DEEP/EMERGENT/MARSHES	X	X			X	X		X	X			X	X	X	X		
	ANABOGA RIVER	SPRING/DEEP/EMERGENT/MARSHES	X	X			X	X		X	X			X	X	X	X		
	ANABOGA RIVER	SPRING/DEEP/EMERGENT/MARSHES	X	X			X	X		X	X			X	X	X	X		
609.12	DEATH VALLEY HYDROLOGIC SUBAREA																		
	MINOR SURFACE WATERS		X	X			X			X	X			X		X			
	MINOR WETLANDS	SPRING/DEEP/EMERGENT/MARSHES	X	X			X	X		X	X			X	X		X	X	
609.13	WINDMILL WASH HYDROLOGIC SUBAREA																		
	MINOR SURFACE WATERS		X	X			X			X	X			X		X			
	MINOR WETLANDS	SPRING/DEEP/EMERGENT/MARSHES	X	X			X	X		X	X			X	X		X	X	
609.20	STAIRMAN WELLS HYDROLOGIC AREA																		
	MINOR SURFACE WATERS		X	X			X			X	X			X		X			
	MINOR WETLANDS	SPRING/DEEP/EMERGENT/MARSHES	X	X			X	X		X	X			X	X		X	X	
609.21	ANABOGA HYDROLOGIC SUBAREA																		
	MINOR SURFACE WATERS		X	X			X			X	X			X		X			
	MINOR WETLANDS	SPRING/DEEP/EMERGENT/MARSHES	X	X			X	X		X	X			X	X		X	X	
609.22	RED PASS HYDROLOGIC SUBAREA																		
	RED PASS LAKE	LAKE	X				X			X	X			X	X	X	X		
	RED PASS LAKE	LAKE	X				X			X	X			X	X	X	X		
	MINOR SURFACE WATERS		X	X			X			X	X			X	X		X	X	
	MINOR WETLANDS	SPRING/DEEP/EMERGENT/MARSHES	X	X			X	X		X	X			X	X		X	X	
609.23	WILLOW HYDROLOGIC SUBAREA																		
	WILLOW LAKE	LAKE	X				X			X	X			X	X	X	X		
	WILLOW LAKE	LAKE	X				X			X	X			X	X	X	X		
	WILLOW LAKE	LAKE	X				X			X	X			X	X	X	X		
	WILLOW LAKE	LAKE	X				X			X	X			X	X	X	X		
	WILLOW LAKE	LAKE	X				X			X	X			X	X	X	X		
609.24	SHADOW HYDROLOGIC SUBAREA																		
	SHADOW LAKE	LAKE	X	X			X			X	X			X	X	X	X		
	SHADOW LAKE	LAKE	X	X			X			X	X			X	X	X	X		
	SHADOW LAKE	LAKE	X	X			X			X	X			X	X	X	X		
609.30	RYAN HYDROLOGIC AREA																		
	MINOR SURFACE WATERS		X	X			X			X	X			X		X			
	MINOR WETLANDS	SPRING/DEEP/EMERGENT/MARSHES	X	X			X	X		X	X			X	X		X	X	
609.31	FUNDACE CREEK HYDROLOGIC SUBAREA																		
	MINOR SURFACE WATERS		X	X			X			X	X			X		X			
	MINOR WETLANDS	SPRING/DEEP/EMERGENT/MARSHES	X	X			X	X		X	X			X	X		X	X	
609.32	GREENWATER HYDROLOGIC SUBAREA																		
	MINOR SURFACE WATERS		X	X			X			X	X			X		X			
	MINOR WETLANDS	SPRING/DEEP/EMERGENT/MARSHES	X	X			X	X		X	X			X	X		X	X	

**TABLE 2-1. BENEFICIAL USES OF SURFACE WATERS OF THE LAHONTAN REGION**

RTU No.	HYDROLOGIC UNIT/SUBUNIT DRAINAGE FEATURE	WATERBODY CLASS MODIFIER	BENEFICIAL USES														RECEIVING WATER		
			WTR	AGRI	IND	DOM	RECR	WILDL	WATER	WATER	WATER	WATER	WATER	WATER	WATER	WATER			
609.40	AMAROGOSA DESERT HYDROLOGIC AREA																		
	MINOR SURFACE WATERS		X	X															AMAROGOSA RIVER
	MINOR WETLANDS		X	X															AMAROGOSA RIVER
609.41	CALICO HYDROLOGIC SUBAREA																		
	MINOR SURFACE WATERS		X	X															AMAROGOSA RIVER
	MINOR WETLANDS		X	X															AMAROGOSA RIVER
609.42	ESCONDIDO HYDROLOGIC SUBAREA																		
	MINOR SURFACE WATERS		X	X															AMAROGOSA RIVER
	MINOR WETLANDS		X	X															AMAROGOSA RIVER
609.43	CHICO HYDROLOGIC SUBAREA																		
	MINOR SURFACE WATERS		X	X															AMAROGOSA RIVER
	MINOR WETLANDS		X	X															AMAROGOSA RIVER
609.44	CALIFORNIA VALLEY HYDROLOGIC SUBAREA																		
	MINOR SURFACE WATERS		X	X															CALIFORNIA VALLEY ON
	MINOR WETLANDS		X	X															CALIFORNIA VALLEY ON
610.00	PANORAMA HYDROLOGIC UNIT																		
	MINOR SURFACE WATERS		X	X															INTERMOUNTAIN VALLEY ON
	MINOR WETLANDS		X	X															INTERMOUNTAIN VALLEY ON
611.00	RESCUITE HYDROLOGIC UNIT																		
	MINOR SURFACE WATERS		X	X															RESCUITE VALLEY ON
	MINOR WETLANDS		X	X															RESCUITE VALLEY ON
612.00	EVANPAH HYDROLOGIC UNIT																		
	MINOR SURFACE WATERS		X	X															EVANPAH VALLEY ON
	MINOR WETLANDS		X	X															EVANPAH VALLEY ON
613.00	UNIONHEAD HYDROLOGIC UNIT																		
	MINOR SURFACE WATERS		X	X															UNIONHEAD VALLEY ON
	MINOR WETLANDS		X	X															UNIONHEAD VALLEY ON
613.10	LOFT LAKE HYDROLOGIC AREA																		
	MINOR SURFACE WATERS		X	X															INTERMOUNTAIN VALLEY ON
	MINOR WETLANDS		X	X															INTERMOUNTAIN VALLEY ON
613.20	ONG LAKE HYDROLOGIC AREA																		
	MINOR SURFACE WATERS		X	X															ONG LAKE
	MINOR WETLANDS		X	X															ONG LAKE
614.00	LEACH HYDROLOGIC UNIT																		
	MINOR SURFACE WATERS		X	X															LEACH VALLEY ON
	MINOR WETLANDS		X	X															LEACH VALLEY ON
615.00	GRANITE HYDROLOGIC UNIT																		
	MINOR SURFACE WATERS		X	X															GRANITE VALLEY ON
	MINOR WETLANDS		X	X															GRANITE VALLEY ON
615.10	DELAKE HYDROLOGIC AREA																		
	MINOR SURFACE WATERS		X	X															DELAKE VALLEY ON
	MINOR WETLANDS		X	X															DELAKE VALLEY ON
615.20	DELAKE HYDROLOGIC AREA																		
	MINOR SURFACE WATERS		X	X															DELAKE VALLEY ON
	MINOR WETLANDS		X	X															DELAKE VALLEY ON
616.00	DELAKE HYDROLOGIC UNIT																		
	MINOR SURFACE WATERS		X	X															DELAKE VALLEY ON
	MINOR WETLANDS		X	X															DELAKE VALLEY ON



**TABLE 2-1. BENEFICIAL USES OF SURFACE WATERS OF THE LAHONTAN REGION**

ITU No.	HYDROLOGIC UNIT/SUBUNIT DRAINAGE FEATURE	WATERBODY CLASS MODIFIER	BENEFICIAL USES																RECEIVING WATER
			WTR	IRRG	IND	IND	IND	IND	IND	IND	IND	IND	IND	IND	IND	IND	IND		
	MINOR WETLANDS	SPRINGS/DEEPS/EMERGENT/MARSHES	X	X	X	X	X	X	X	X	X	X	X	X	X	X			
617.00	<b>GOLDSTONE HYDROLOGIC UNIT</b>																		
	GOLDSTONE LAKE	ALTA 1 LAKE	X		X			X	X	X	X	X	X				INTERIALLY DRAINED LAKE		
	SPRING LAKE	ALTA 1 LAKE	X		X			X	X	X	X	X	X				INTERIALLY DRAINED LAKE		
	GOLDSTONE LAKE	LAKE	X		X			X	X	X	X	X	X						
	MINOR SURFACE WATERS		X	X	X	X	X	X	X	X	X	X	X	X					
	MINOR WETLANDS	SPRINGS/DEEPS/EMERGENT/MARSHES	X	X	X	X	X	X	X	X	X	X	X			X	X		
618.00	<b>COYOTE HYDROLOGIC UNIT</b>																		
	BARNEY SPRING	SPRINGS/DEEPS/EMERGENT/MARSHES	X	X	X	X	X	X	X	X	X	X	X			X	COYOTE LAKE ON		
	JACK SPRING	SPRINGS	X	X	X	X	X	X	X	X	X	X	X				COYOTE LAKE ON		
	COYOTE LAKE		X		X	X	X	X	X	X	X	X	X				COYOTE LAKE		
	JACK RABBIT SPRING		X		X	X	X	X	X	X	X	X	X				COYOTE LAKE		
	MINOR SURFACE WATERS		X	X	X	X	X	X	X	X	X	X	X						
	MINOR WETLANDS	SPRINGS/DEEPS/EMERGENT/MARSHES	X	X	X	X	X	X	X	X	X	X	X			X	X		
619.00	<b>SUFFERCORN HYDROLOGIC UNIT</b>																		
	SUFFERCORN LAKE	LAKE	X					X	X	X	X	X	X				SUFFERCORN LAKE		
	COLEMAN SPRING	SPRINGS	X	X				X	X	X	X	X	X				SUFFERCORN LAKE		
	EDWARDS LAKE	LAKE	X					X	X	X	X	X	X				SUFFERCORN LAKE		
	MINOR SURFACE WATERS		X	X	X	X	X	X	X	X	X	X	X						
	MINOR WETLANDS	SPRINGS/DEEPS/EMERGENT/MARSHES	X	X	X	X	X	X	X	X	X	X	X			X	X		
620.00	<b>BALLAST HYDROLOGIC UNIT</b>																		
	MINOR SURFACE WATERS		X	X	X	X	X	X	X	X	X	X	X						
	MINOR WETLANDS	SPRINGS/DEEPS/EMERGENT/MARSHES	X	X	X	X	X	X	X	X	X	X	X			X	X		
620.10	<b>WINDMILL HYDROLOGIC AREA</b>																		
	MINOR SURFACE WATERS		X		X			X	X	X	X	X	X						
	MINOR WETLANDS	SPRINGS/DEEPS/EMERGENT/MARSHES	X	X	X	X	X	X	X	X	X	X	X			X	X		
620.20	<b>MILBURN HYDROLOGIC AREA</b>																		
	MINOR SURFACE WATERS		X	X	X	X	X	X	X	X	X	X	X	X					
	MINOR WETLANDS	SPRINGS/DEEPS/EMERGENT/MARSHES	X	X	X	X	X	X	X	X	X	X	X			X	X		
620.21	<b>WHITE ROCK HYDROLOGIC SUBAREA</b>																		
	MINOR SURFACE WATERS		X	X	X	X	X	X	X	X	X	X	X						
	MINOR WETLANDS	SPRINGS/DEEPS/EMERGENT/MARSHES	X	X	X	X	X	X	X	X	X	X	X			X	X		
620.22	<b>WILD ROSE PEAK HYDROLOGIC SUBAREA</b>																		
	MINOR SURFACE WATERS		X	X	X	X	X	X	X	X	X	X	X						
	MINOR WETLANDS	SPRINGS/DEEPS/EMERGENT/MARSHES	X	X	X	X	X	X	X	X	X	X	X			X	X		
620.30	<b>LEE FLAT HYDROLOGIC AREA</b>																		
	MINOR SURFACE WATERS		X		X	X	X	X	X	X	X	X	X						
	MINOR WETLANDS	SPRINGS/DEEPS/EMERGENT/MARSHES	X	X	X	X	X	X	X	X	X	X	X			X	X		
620.40	<b>SANTA ROSA FLAT HYDROLOGIC AREA</b>																		
	MINOR SURFACE WATERS		X		X	X	X	X	X	X	X	X	X						
	MINOR WETLANDS	SPRINGS/DEEPS/EMERGENT/MARSHES	X	X	X	X	X	X	X	X	X	X	X			X	X		
620.41	<b>WOLFPEAK MEAD HYDROLOGIC SUBAREA</b>																		
	MINOR SURFACE WATERS		X	X	X	X	X	X	X	X	X	X	X						
	MINOR WETLANDS	SPRINGS/DEEPS/EMERGENT/MARSHES	X	X	X	X	X	X	X	X	X	X	X			X	X		
620.42	<b>RAIMON HYDROLOGIC SUBAREA</b>																		
	MINOR SURFACE WATERS		X		X	X	X	X	X	X	X	X	X						
	MINOR WETLANDS	SPRINGS/DEEPS/EMERGENT/MARSHES	X	X	X	X	X	X	X	X	X	X	X			X	X		
620.43	<b>STEVEN DOLLAR HYDROLOGIC SUBAREA</b>																		
	MINOR SURFACE WATERS		X		X	X	X	X	X	X	X	X	X						
	MINOR WETLANDS	SPRINGS/DEEPS/EMERGENT/MARSHES	X	X	X	X	X	X	X	X	X	X	X			X	X		
620.50	<b>DARWIN HYDROLOGIC AREA</b>																		
	MINOR SURFACE WATERS		X	X	X	X	X	X	X	X	X	X	X						
	MINOR WETLANDS	SPRINGS/DEEPS/EMERGENT/MARSHES	X	X	X	X	X	X	X	X	X	X	X			X	X		
620.60	<b>PANAMINT VALLEY HYDROLOGIC AREA</b>																		
	HELANDS SPRING, DOAN TILL POND	SPRING, CREEK	X		X			X	X	X	X	X	X				PANAMINT VALLEY ON		
	REDBOOKS SPRING	SPRING	X		X			X	X	X	X	X	X				PANAMINT VALLEY ON		
	GOLEER CREEK (UNION) (REAR)	SPRING	X		X			X	X	X	X	X	X				PANAMINT VALLEY ON		
	MINOR SURFACE WATERS		X	X	X	X	X	X	X	X	X	X	X						
	MINOR WETLANDS	SPRINGS/DEEPS/EMERGENT/MARSHES	X	X	X	X	X	X	X	X	X	X	X			X	X		
620.70	<b>BROWN HYDROLOGIC AREA</b>																		
	MINOR SURFACE WATERS		X		X	X	X	X	X	X	X	X	X						
	MINOR WETLANDS	SPRINGS/DEEPS/EMERGENT/MARSHES	X	X	X	X	X	X	X	X	X	X	X			X	X		
620.80	<b>ROBBERS HYDROLOGIC AREA</b>																		
	LEAD POND SPRING	SPRING	X		X			X	X	X	X	X	X				PICT ROCK SPR., PANAMINT MOUNT.		
	MINOR SURFACE WATERS		X		X			X	X	X	X	X	X						
	MINOR WETLANDS		X	X	X	X	X	X	X	X	X	X	X			X	X		
621.00	<b>TRONA HYDROLOGIC UNIT</b>																		
	CLASSED, WY LAKE PFT.	SALT FLAT LAKE		X	X			X	X			X	X				TERMINAL DRAINED LAKE		
	MINOR SURFACE WATERS		X		X	X	X	X	X	X	X	X	X						
	MINOR WETLANDS		X	X	X	X	X	X	X	X	X	X	X			X	X		
621.10	<b>CHARLES VALLEY HYDROLOGIC AREA</b>																		
	PEACH SPRINGS	SPRING	X		X			X	X	X	X	X	X				CHARLES VALLEY DRAINAGE		
	FRANKS SPRING (ON THE NE CORNER OF TRONA W. QUAD)	SPRING	X		X			X	X	X	X	X	X				CHARLES VALLEY ON		
	SPRINGS ON THE W. CORNER OF CANYON	SPRING	X		X			X	X	X	X	X	X				CHARLES VALLEY ON		
	MINOR SURFACE WATERS		X		X			X	X	X	X	X	X						
	MINOR WETLANDS	SPRING	X	X	X	X	X	X	X	X	X	X	X			X	X		
621.20	<b>SALT WELLS HYDROLOGIC AREA</b>																		
	MINOR SURFACE WATERS		X		X			X	X	X	X	X	X						
	MINOR WETLANDS		X	X	X	X	X	X	X	X	X	X	X			X	X		
621.30	<b>PICT ROCK HYDROLOGIC AREA</b>																		
	PICT SPRING	SPRING	X		X			X	X	X	X	X	X				CHARLES VALLEY		
	CHARLES VALLEY SPRING	SPRING	X		X			X	X	X	X	X	X				CHARLES VALLEY		

**TABLE 2-1. BENEFICIAL USES OF SURFACE WATERS OF THE LAHONTAN REGION**

W.D. No.	HYDROLOGIC UNIT/SUBUNIT DRAINAGE FEATURE	WATERBODY CLASS MODIFIER	BENEFICIAL USES														RECEIVING WATER			
			WTR	IRR	PAO	IND	WPL	WRE	WAV	WOM	WOC-1	WOC-2	WOC-3	WOC-4	WOC-5	WOC-6		WOC-7	WOC-8	WOC-9
	MUDW. S. P. H. A. S. W. A. S. S. E. S.		X																	
	MUDW. W. A. S. S. E. S.	WETLANDS	X																X	X
622.00	COAD HYDROLOGIC UNIT																			
	WATER RESERVE WATERS		X	X		X				X	X	X								
	MUDW. W. A. S. S. E. S.		X	X		X				X	X	X							X	X
622.10	MUDW. W. A. S. S. E. S.																			
	WATER RESERVE WATERS		X	X		X				X	X	X								
	MUDW. W. A. S. S. E. S.		X	X		X				X	X	X							X	X
622.20	WATER RESERVE WATERS		X	X		X				X	X	X								
	MUDW. W. A. S. S. E. S.	WETLANDS	X							X	X	X							X	X
623.00	UPPER CANYON HYDROLOGIC UNIT																			
	WATER RESERVE WATERS		X	X		X				X	X	X								
	MUDW. W. A. S. S. E. S.	WETLANDS	X	X		X				X	X	X							X	X
624.00	INDIAN WELLS HYDROLOGIC UNIT																			
	WATER RESERVE WATERS		X	X		X				X	X	X								
	MUDW. W. A. S. S. E. S.		X	X		X				X	X	X								
	MUDW. W. A. S. S. E. S.	WETLANDS	X	X		X				X	X	X							X	X
624.10	BORE HYDROLOGIC AREA																			
	LITTLE LAKE	LAKE	X	X		X				X	X	X								
	LITTLE LAKE CANYON LAKE		X	X		X				X	X	X								
	INTERMEDIATE CANYON		X	X		X				X	X	X							X	
	MUDW. S. P. H. A. S. W. A. S. S. E. S.		X	X		X				X	X	X							X	
	MUDW. W. A. S. S. E. S.	WETLANDS	X	X		X				X	X	X							X	X
624.20	INDIAN WELLS HYDROLOGIC AREA																			
	INDIAN WELLS CANYON CREEK	INTERMEDIATE CANYON	X	X		X				X	X	X								
	LAKE DEEP CANYON	LAKE	X	X		X				X	X	X								
	LAKE DEEP	SPRING	X	X		X				X	X	X							X	
624.20	INDIAN WELLS (continued)																			
	INDIAN WELLS CANYON CREEK	SPRING	X	X		X				X	X	X								
	INDIAN WELLS CANYON CREEK	SPRING	X	X		X				X	X	X								
	INDIAN WELLS CANYON CREEK	SPRING	X	X		X				X	X	X								
	INDIAN WELLS CANYON CREEK	SPRING	X	X		X				X	X	X								
	INDIAN WELLS CANYON CREEK	SPRING	X	X		X				X	X	X								
	INDIAN WELLS CANYON CREEK	SPRING	X	X		X				X	X	X								
	INDIAN WELLS CANYON CREEK	SPRING	X	X		X				X	X	X								
	INDIAN WELLS CANYON CREEK	SPRING	X	X		X				X	X	X								
	INDIAN WELLS CANYON CREEK	SPRING	X	X		X				X	X	X								
	INDIAN WELLS CANYON CREEK	SPRING	X	X		X				X	X	X								
	INDIAN WELLS CANYON CREEK	SPRING	X	X		X				X	X	X								
	INDIAN WELLS CANYON CREEK	SPRING	X	X		X				X	X	X								
	INDIAN WELLS CANYON CREEK	SPRING	X	X		X				X	X	X								
	INDIAN WELLS CANYON CREEK	SPRING	X	X		X				X	X	X								
	INDIAN WELLS CANYON CREEK	SPRING	X	X		X				X	X	X								
	INDIAN WELLS CANYON CREEK	SPRING	X	X		X				X	X	X								
	INDIAN WELLS CANYON CREEK	SPRING	X	X		X				X	X	X								
	INDIAN WELLS CANYON CREEK	SPRING	X	X		X				X	X	X								
	INDIAN WELLS CANYON CREEK	SPRING	X	X		X				X	X	X								
	INDIAN WELLS CANYON CREEK	SPRING	X	X		X				X	X	X								
	INDIAN WELLS CANYON CREEK	SPRING	X	X		X				X	X	X								
	INDIAN WELLS CANYON CREEK	SPRING	X	X		X				X	X	X								
	INDIAN WELLS CANYON CREEK	SPRING	X	X		X				X	X	X								
	INDIAN WELLS CANYON CREEK	SPRING	X	X		X				X	X	X								
	INDIAN WELLS CANYON CREEK	SPRING	X	X		X				X	X	X								
	INDIAN WELLS CANYON CREEK	SPRING	X	X		X				X	X	X								
	INDIAN WELLS CANYON CREEK	SPRING	X	X		X				X	X	X								
	INDIAN WELLS CANYON CREEK	SPRING	X	X		X				X	X	X								
	INDIAN WELLS CANYON CREEK	SPRING	X	X		X				X	X	X								
	INDIAN WELLS CANYON CREEK	SPRING	X	X		X				X	X	X								
	INDIAN WELLS CANYON CREEK	SPRING	X	X		X				X	X	X								
	INDIAN WELLS CANYON CREEK	SPRING	X	X		X				X	X	X								

**TABLE 2-1. BENEFICIAL USES OF SURFACE WATERS OF THE LAHONTAN REGION**

[illegible]

**TABLE 2-1. BENEFICIAL USES OF SURFACE WATERS OF THE LAHONTAN REGION**

HU No.	HYDROLOGIC UNIT/SUBUNIT DRAINAGE FEATURE	WATERBODY CLASS MODIFIER	BENEFICIAL USES																RECEIVING WATER
			WTR	AGR	IND	WTR	WTR	WTR	WTR	WTR	WTR	WTR	WTR	WTR	WTR	WTR	WTR		
628.41	GRASS VALLEY HYDROLOGIC SUBAREA																		
	MINOR SURFACE WATER		X	X															
	MINOR WETLANDS	WETLANDS	X	X		X	X		X	X		X	X		X	X			
628.42	HARPER VALLEY HYDROLOGIC SUBAREA																		
	SPRING	SPRING	X	X		X			X	X		X	X		X			HARPER VALLEY (GROUNDWATER)	
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			INTERNALLY DRAINAGE LAKE	
	UPPER SPRING	SPRING													X				
	HARPER LAKE WETLANDS	WETLANDS	X	X		X			X	X		X	X		X	X		HARPER LAKE	
	MINOR SURFACE WATER		X	X		X			X	X		X	X		X				
628.50	LOWER MONTANA HYDROLOGIC AREA																		
	MINOR SURFACE WATER		X	X		X			X	X		X	X		X				
	MINOR WETLANDS	WETLANDS	X	X		X	X		X	X		X	X		X	X			
628.60	NEMESSY SPRINGS HYDROLOGIC AREA																		
	MINOR SURFACE WATER		X	X		X			X	X		X	X		X				
	MINOR WETLANDS	WETLANDS	X	X		X	X		X	X		X	X		X	X			
628.61	FAIR HOLE HYDROLOGIC SUBAREA																		
	MINOR SURFACE WATER		X	X		X			X	X		X	X		X				
	MINOR WETLANDS	WETLANDS	X	X		X	X		X	X		X	X		X	X			
628.62	TRACY VALLEY HYDROLOGIC SUBAREA																		
	MINOR SURFACE WATER		X	X		X			X	X		X	X		X				
	MINOR WETLANDS	WETLANDS	X	X		X	X		X	X		X	X		X	X			
628.70	AYTON HYDROLOGIC AREA																		
	MINOR SURFACE WATER		X	X		X			X	X		X	X		X				
	MINOR WETLANDS	WETLANDS	X	X		X	X		X	X		X	X		X	X			
628.71	DAVIS HYDROLOGIC SUBAREA																		
	MINOR SURFACE WATER		X	X		X			X	X		X	X		X			MONTANA A. FORD RESERVOIR	
	MINOR SURFACE WATER		X	X		X			X	X		X	X		X				
	MINOR WETLANDS	WETLANDS	X	X		X	X		X	X		X	X		X	X			
628.72	CHICKSEE HYDROLOGIC SUBAREA																		
	UPPER SPRING	WETLANDS	X	X		X			X	X		X	X		X			CHICKSEE VALLEY LSP	
	CHICKSEE LAKE (LARGE AND SMALL)	WETLANDS	X	X		X			X	X		X	X		X			CHICKSEE LAKE	
	MINOR SURFACE WATER		X	X		X			X	X		X	X		X				
	MINOR WETLANDS	WETLANDS	X	X		X	X		X	X		X	X		X	X			
628.73	LANDFORD HYDROLOGIC SUBAREA																		
	MINOR SURFACE WATER		X	X		X			X	X		X	X		X				
	MINOR SURFACE WATER	WETLANDS	X	X		X	X		X	X		X	X		X	X			
628.86	HAKER HYDROLOGIC AREA																		
	MINOR SURFACE WATER		X	X		X			X	X		X	X		X				
	MINOR WETLANDS	WETLANDS	X	X		X	X		X	X		X	X		X	X			
628.81	STIVERS LAKE HYDROLOGIC SUBAREA																		
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			STIVERS LAKE (STIVERS LAKE BASIN)	
	SHALLOW LAKE (WETLANDS)	WETLANDS	X	X		X			X	X		X	X		X			STIVERS LAKE	
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			STIVERS LAKE	
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			STIVERS LAKE	
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			STIVERS LAKE	
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			STIVERS LAKE	
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			STIVERS LAKE	
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			STIVERS LAKE	
MINOR SURFACE WATER		X	X		X			X	X		X	X		X					
628.81	MINOR WETLANDS	WETLANDS	X	X		X	X		X	X		X	X		X	X			
	MINOR SURFACE WATER		X	X		X			X	X		X	X		X				
	MINOR WETLANDS	WETLANDS	X	X		X	X		X	X		X	X		X	X			
628.82	BLAIR LAKE HYDROLOGIC SUBAREA																		
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			INTERNALLY DRAINAGE LAKE	
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			INTERNALLY DRAINAGE LAKE	
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			INTERNALLY DRAINAGE LAKE	
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			INTERNALLY DRAINAGE LAKE	
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			INTERNALLY DRAINAGE LAKE	
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			INTERNALLY DRAINAGE LAKE	
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			INTERNALLY DRAINAGE LAKE	
MINOR SURFACE WATER		X	X		X			X	X		X	X		X					
628.82	MINOR WETLANDS	WETLANDS	X	X		X	X		X	X		X	X		X	X			
	MINOR SURFACE WATER		X	X		X			X	X		X	X		X				
	MINOR WETLANDS	WETLANDS	X	X		X	X		X	X		X	X		X	X			
628.90	BLAIR LAKE HYDROLOGIC AREA																		
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			SHALLOW LAKE	
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			SHALLOW LAKE	
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			SHALLOW LAKE	
628.90	BLAIR LAKE HYDROLOGIC AREA																		
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			SHALLOW LAKE	
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			SHALLOW LAKE	
628.90	BLAIR LAKE HYDROLOGIC AREA																		
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			SHALLOW LAKE	
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			SHALLOW LAKE	
628.90	BLAIR LAKE HYDROLOGIC AREA																		
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			SHALLOW LAKE	
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			SHALLOW LAKE	
628.90	BLAIR LAKE HYDROLOGIC AREA																		
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			SHALLOW LAKE	
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			SHALLOW LAKE	
628.90	BLAIR LAKE HYDROLOGIC AREA																		
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			SHALLOW LAKE	
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			SHALLOW LAKE	
628.90	BLAIR LAKE HYDROLOGIC AREA																		
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			SHALLOW LAKE	
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			SHALLOW LAKE	
628.90	BLAIR LAKE HYDROLOGIC AREA																		
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			SHALLOW LAKE	
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			SHALLOW LAKE	
628.90	BLAIR LAKE HYDROLOGIC AREA																		
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			SHALLOW LAKE	
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			SHALLOW LAKE	
628.90	BLAIR LAKE HYDROLOGIC AREA																		
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			SHALLOW LAKE	
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			SHALLOW LAKE	
628.90	BLAIR LAKE HYDROLOGIC AREA																		
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			SHALLOW LAKE	
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			SHALLOW LAKE	
628.90	BLAIR LAKE HYDROLOGIC AREA																		
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			SHALLOW LAKE	
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			SHALLOW LAKE	
628.90	BLAIR LAKE HYDROLOGIC AREA																		
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			SHALLOW LAKE	
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			SHALLOW LAKE	
628.90	BLAIR LAKE HYDROLOGIC AREA																		
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			SHALLOW LAKE	
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			SHALLOW LAKE	
628.90	BLAIR LAKE HYDROLOGIC AREA																		
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			SHALLOW LAKE	
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			SHALLOW LAKE	
628.90	BLAIR LAKE HYDROLOGIC AREA																		
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			SHALLOW LAKE	
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			SHALLOW LAKE	
628.90	BLAIR LAKE HYDROLOGIC AREA																		
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			SHALLOW LAKE	
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			SHALLOW LAKE	
628.90	BLAIR LAKE HYDROLOGIC AREA																		
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			SHALLOW LAKE	
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			SHALLOW LAKE	
628.90	BLAIR LAKE HYDROLOGIC AREA																		
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			SHALLOW LAKE	
	SHALLOW LAKE	SHALLOW LAKE	X	X		X			X	X		X	X		X			SHALLOW LAKE	
628.90	BLAIR LAKE HYDROLOGIC AREA																		
	SHALLOW LAKE	SHALLOW LAKE	X	X		X</													

**Changes to Table 2-1 of the Lahontan Basin Plan made in plan amendments adopted in July 2000 and finally approved in April 2002.** The table below shows the beneficial uses remaining for these water after removal of the potential Municipal and Domestic Supply (MUN) use designation. Rows for individual water bodies in the Table below will be revised in or added to Table 2-1 of the Basin Plan when the plan is reprinted; fonts will be revised to match the rest of the table. See Chapter 2 of the Basin Plan for the meaning of abbreviations in the “Beneficial Use” columns.

HU No.	HYDRO-LOGIC UNIT/ SUBUNIT DRAINAGE FEATURE	WATERBODY CLASS MODIFIER	BENEFICIAL USES																										Receiving Water
			M U N	A G R	P R O	I N D	G W	F R S H	N A V	P O W	R E C - 1	R E C - 2	C O C - M	A Q U I F E R	C O A L D	S A L I D	W I D	B I O R E	R A R E	M I G R N	S P W	W Q E	F L D						
637.20	Wendel Hot Springs	Hot Springs	x				x	x		x	x																	Honey Lake	
637.20	Amedee Hot Springs	Hot Springs	x				x	x			x	x																Honey Lake	
631.40	Hot Creek	Perennial Stream	x				x				x	x																Little Walker River	
631.40	Fales Hot Springs	Springs	x				x				x	x																Hot Creek	
603.10	Little Hot Creek	Perennial Stream	x				x	x			x	x																Hot Creek or Owens River	
603.10	Little Alkali Lake	Alkali Lake	x				x	x			x	x																Crowley Lake	
603.20	Keough Hot Springs	Springs	x				x				x	x																Owens River	
605.00	Deep Springs Lake	Intermittent Lake					x				x	x																Deep Springs Val. GW	
609.00	Amargosa River	Intermittent Stream	x				x				x	x																Amargosa Subarea GW	

**Summary of Beneficial Uses for Waterbodies Identified in the Nevada Administrative Code**

NAC	Name	Description	Beneficial Uses								Aquatic species of concern			
			IRR	STOCK	REC-1	REC-2	IND	MUN	WILD	AQUATIC		AESTHETIC	ENHANCE	MARSH
CLASS A WATERS														
445A.124	Various waterbodies		X	X	X	X			A	X	X			
CLASS B WATERS														
445A.125	Various waterbodies		X	X	X	X	X		B	X	X			
CLASS C WATERS														
445A.126	Various waterbodies		X	X	X	X	X	X	C	X	X			
CLASS D WATERS														
445A.127	Various waterbodies		X	X			X	X		X	X			
CARSON RIVER BASIN														
445A.147	West Fork Carson River	At stateline	X	X	X	X	X	X	X	X	X			rainbow trout, brown trout
445A.148	Bryant Creek	At stateline	X	X	X	X	X	X	X	X	X			rainbow trout, brown trout
445A.149	East Fork Carson River	At stateline	X	X	X	X	X	X	X	X	X			rainbow trout, brown trout
445A.150	East Fork Carson River	Stateline to Highway 395	X	X	X	X	X	X	X	X	X			rainbow trout, brown trout
445A.151	East Fork Carson River	Highway 395 to Muller Lane	X	X	X	X	X	X	X	X	X			rainbow trout, brown trout
445A.152	EF/WF & Carson River	EF at Muller to Genoa Lane & WF at stateline to Genoa Lane	X	X	X	X	X	X	X	X	X			catfish, rainbow trout, brown trout
445A.153	Carson River	Genoa Lane to Cradlebaugh Bridge	X	X	X	X	X	X	X	X	X			catfish, rainbow trout, brown trout
445A.154	Carson River	Cradlebaugh Bridge to Mexican Ditch Gage	X	X	X	X	X	X	X	X	X			rainbow trout, brown trout
445A.155	Carson River	Mexican Ditch Gage to New Empire	X	X	X	X	X	X	X	X	X			smallmouth bass, rainbow trout, brown trout
445A.156	Carson River	New Empire to Dayton Bridge	X	X	X	X	X	X	X	X	X			walleye, channel catfish, white bass
445A.157	Carson River	Dayton Bridge to Weeks Bridge	X	X	X	X	X	X	X	X	X			walleye, channel catfish, white bass
445A.158	Carson River	Weeks Bridge to Lahontan Dam	X	X	X	X	X	X	X	X	X			walleye, channel catfish, white bass
WALKER RIVER BASIN														
445A.160	West Walker River	At stateline	X	X	X	X	X	X	X	X	X			mountain whitefish, rainbow trout, brown trout
445A.161	Topaz Lake	Various points in Topaz Lake	X	X	X	X	X	X	X	X	X			rainbow trout, brown trout, cutthroat trout, kokone salmon, silver salmon
445A.162	West Walker River	CA stateline to Wellington	X	X	X	X	X	X	X	X	X			mountain whitefish, rainbow trout, brown trout
445A.163	West Walker River	Wellington to confluence with East Walker River	X	X	X	X	X	X	X	X	X			rainbow trout, brown trout
445A.164	Sweetwater Creek	CA stateline to the East Walker River	X	X	X	X	X	X	X	X	X			mountain whitefish, brown trout, brook trout, rainbow trout
445A.165	East Walker River	At stateline	X	X	X	X	X	X	X	X	X			mountain white fish, rainbow trout, brown trout
445A.1655	East Walker River	Stateline to Bridge B-1475	X	X	X	X	X	X	X	X	X			mountain white fish, rainbow trout, brown trout
445A.166	East Walker River	Bridge B-1475 to East/West Walker confluence	X	X	X	X	X	X	X	X	X			brown trout, rainbow trout
445A.167	Walker River	East/West Walker confluence to Weber Reservoir Inlet	X	X	X	X	X	X	X	X	X			channel catfish, largemouth bass
445A.168	Walker River	Weber Reservoir to Walker Lake												
445A.169	Desert Creek	CA stateline to the West Walker River	X	X	X	X	X	X	X	X	X			brown trout, brook trout, rainbow trout

State standards do not apply on tribal lands

Summary of Beneficial Uses for Waterbodies Identified in the Nevada Administrative Code

NAC	Name	Description	Beneficial Uses								Aquatic species of concern		
			IRR	STOCK	REC-1	REC-2	IND	MUN	WILD	AQUATIC		AESTHETIC	ENHANCE
445A.1696	Walker Lake	Walker Lake			X	X			X	X			tui chub, Tahoe sucker, and adult and juvenile Lahontan cutthroat trout
CENTRAL REGION													
445A.171	Chiatovich Creek	Above highway maintenance station	X	X	X	X	X	X	X	X			
445A.172	Indian Creek	Above center of Section 9, T2S, R34E	X	X	X	X	X	X	X	X			
445A.173	Leidy Creek	Above hydroelectric plant	X	X	X	X	X	X	X	X			
COLORADO RIVER BASIN													
445A.175	Virgin River	AZ stateline to Mesquite	X	X		X	X	X		X	X		
445A.176	Virgin River	At stateline	X	X		X	X	X		X	X		
445A.177	Virgin River	Mesquite to river mouth at Lake Mead	X	X		X	X	X		X	X		
445A.178	Beaver Dam Wash	Above Schroeder Reservoir	X	X	X	X	X	X	X	X	X		
445A.192	Colorado River	Lake Mohave Inlet to CA stateline	X	X	X	X	X	X	X	X	X		
445A.193	Colorado River	Hoover Dam to Lake Mohave Inlet	X	X	X	X	X	X	X	X	X		
445A.195	Lake Mead	Excluding area covered by NAC 445A.197	X	X	X	X	X	X	X	X	X		warmwater fishery
445A.197	Lake Mead	West boundary of Las Vegas Bay campground to confluence of Las Vegas Wash	X	X		X	X	X		X	X		warmwater fishery
445A.199	Las Vegas Wash	Confluence of discharges from City of Las Vegas and Clark County wastewater treatment plants to Telephone Road	X	X		X			X	X		X	excluding fish
445A.201	Las Vegas Wash	Telephone Road to Lake Mead	X	X		X				X	X	X	excluding fish
445A.210	Muddy River	River source to Glendale	X	X		X	X	X		X	X		
445A.211	Muddy River	Glendale to Lake Mead	X	X		X	X	X		X	X		
445A.212	Meadow Valley Wash	Bridge above Rox to Muddy River	X	X		X	X	X		X	X		
GREAT SALT LAKE BASIN													
445A.179	Snake Creek	Above fish hatchery	X	X	X	X	X	X	X	X	X		
WESTERN REGION													
445A.180	Smoke Creek	No description given	No beneficial uses defined										
TRUCKEE RIVER BASIN													
445A.181	Bronco Creek	No description given	No beneficial uses defined										
445A.182	Gray Creek	No description given	No beneficial uses defined										
445A.184	Truckee River	At stateline	X	X	X	X	X	X	X	X	X		all life stages of mountain whitefish, rainbow trout, brown trout
445A.185	Truckee River	CA stateline to Idlewild	X	X	X	X	X	X	X	X	X		all life stages of mountain whitefish, rainbow trout, brown trout
445A.186	Truckee River	Idlewild to East McCarran Blvd	X	X	X	X	X	X	X	X	X		juvenile and adult rainbow trout, juvenile and adult brown trout
445A.187	Truckee River	East McCarran Blvd to Lockwood	X	X	X	X	X	X	X	X	X		juvenile and adult rainbow trout, juvenile and adult brown trout
445A.188	Truckee River	Lockwood to Derby Dam	X	X	X	X	X	X	X	X	X		early spawning Lahontan cutthroat trout and their incubation, larvae, juveniles and migration
445A.189	Truckee River	Derby Dam to Wadsworth	X	X	X	X	X	X	X	X	X		

# Summary of Beneficial Uses for Waterbodies Identified in the Nevada Administrative Code

NAC	Name	Description	Beneficial Uses								Aquatic species of concern				
			IRR	STOCK	REC-1	REC-2	IND	MUN	WILD	AQUATIC		AESTHETIC	ENHANCE	MARSH	
445A.190	Truckee River	Wadsworth to Pyramid Lake													
445A.191	Lake Tahoe	Lake Tahoe	X	X	X	X	X	X	X	X	X	X		coldwater fishery	
445A.1915	Tributaries to Lake Tahoe	All tributaries	X	X	X	X	X	X	X	X	X	X	X	coldwater fishery	
HUMBOLDT RIVER BASIN															
445A.203	Humboldt River	Source to Osino	X	X	X	X	X	X	X	X	X	X		warmwater fishery	
445A.204	Humboldt River	Osino to Palisade	X	X	X	X	X	X	X	X	X	X		warmwater fishery	
445A.205	Humboldt River	Palisade to Battle Mountain	X	X	X	X	X	X	X	X	X	X		warmwater fishery	
445A.206	Humboldt River	Battle Mountain to State Highway 789	X	X	X	X	X	X	X	X	X	X		warmwater fishery	
445A.207	Humboldt River	Cornus to Imlay	X	X	X	X	X	X	X	X	X	X		warmwater fishery	
445A.208	Humboldt River	Imlay to Woosley	X	X	X	X	X	X	X	X	X	X		warmwater fishery	
SNAKE RIVER BASIN															
445A.215	Big Goose Creek	Entire reach	X	X	X	X	X	X	X	X	X	X			
445A.216	Salmon Falls Creek	Entire reach	X	X	X	X	X	X	X	X	X	X			
445A.217	Shoshone Creek	Entire reach	X	X	X	X	X	X	X	X	X	X			
445A.218	EF Jarbridge River	Entire reach	X	X	X	X	X	X	X	X	X	X			
445A.219	Jarbridge River	Upstream from Jarbridge	X	X	X	X	X	X	X	X	X	X			
445A.220	Jarbridge River	Jarbridge to stateline	X	X	X	X	X	X	X	X	X	X			
445A.221	West Fork Bruneau River	Entire reach	X	X	X	X	X	X	X	X	X	X			
445A.222	East Fork Owyhee River	Wildhorse Reservoir to Mill Creek	X	X	X	X	X	X	X	X	X	X			
445A.223	East Fork Owyhee River	Mill Creek to New China Dam	X	X	X	X	X	X	X	X	X	X			
445A.224	East Fork Owyhee River	New China Dam to stateline	X	X	X	X	X	X	X	X	X	X			
445A.225	South Fork Owyhee River	Entire reach	X	X	X	X	X	X	X	X	X	X			
State standards do not apply on tribal lands															

<b>IRR</b>	Irrigation	<b>WILD</b>	Propagation of wildlife
<b>STOCK</b>	Watering of livestock	<b>AQUATIC</b>	Propagation of aquatic life
<b>REC-1</b>	Recreation involving contact with the water	<b>AESTHETIC</b>	Waters of extraordinary ecological or aesthetic value
<b>REC-2</b>	Recreation not involving contact with the water	<b>ENHANCE</b>	Enhancement of water quality
<b>IND</b>	Industrial supply	<b>MARSH</b>	Maintenance of a freshwater marsh
<b>MUN</b>	Municipal or domestic supply, or both		
<b>A</b>	With treatment by disinfection only		
<b>B</b>	With treatment by disinfection and filtration only		
<b>C</b>	With complete treatment		